# antibodies - online.com







## anti-TLR1 antibody (AA 350-404)





$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μg
Target:	TLR1
Binding Specificity:	AA 350-404
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TLR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Flow Cytometry (FACS), Immunocytochemistry (ICC)

#### **Product Details**

Brand:	Picoband™
Immunogen:	E. coli-derived human TLR1 recombinant protein (Position: F350-D404).
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	Rabbit IgG polyclonal antibody for TLR1 detection. Tested with WB, IHC-P, IHC-F, ICC, FCM, Direct ELISA in Human, Mouse, Rat.

### **Target Details**

Target:	TLR1
Alternative Name:	TLR1 (TLR1 Products)

Target Details	
Background:	Synonyms: Toll-like receptor 1, Toll/interleukin-1 receptor-like protein, TIL, CD281, TLR1, KIAA0012
	Tissue Specificity: Ubiquitous. Highly expressed in spleen, ovary, peripheral blood leukocytes,
	thymus and small intestine.
	Background: Toll-like receptor 1 (TLR1), also called TIL or CD281 is a member of the Toll-like
	receptor family (TLR) of pattern recognition receptors of the innate immune system. This gene
	is mapped to 4p14 by fluorescence in situ hybridization. TLRs are highly conserved from
	Drosophila to humans and share structural and functional similarities. They recognize
	pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and
	mediate the production of cytokines necessary for the development of effective immunity. The
	various TLRs exhibit different patterns of expression. This gene is ubiquitously expressed, and
	at higher levels than other TLR genes. Different length transcripts presumably resulting from
	use of alternative polyadenylation site, and/or from alternative splicing, have been noted for this
	gene.
UniProt:	Q15399
Pathways:	TLR Signaling, Activation of Innate immune Response, Cellular Response to Molecule of
	Bacterial Origin, Toll-Like Receptors Cascades
Application Details	
Application Notes:	Recommended Detection Systems: Enhanced Chemiluminescent Kit with anti-Rabbit IgG
	(ABIN921124) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit
	(SV0002-1) for IHC(P), IHC(F) and ICC.
	Application Details: Western blot, 0.1-0.5 μg/mL
	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/mL
	Immunohistochemistry(Frozen Section), 0.5-1 μg/mL
	Immunocytochemistry, 0.5-1 μg/mL
	Flow Cytometry, 1-3 µg/1x10 <sup>6</sup> cells
	Direct ELISA, 0.1-0.5 μg/mL
Restrictions:	For Research Use only

### Handling


Format: Lyophilized

Reconstitution: Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.

#### Handling


Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05 mg NaN <sub>3</sub> .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month.  It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.


#### **Images**



#### **Western Blotting**

Image 1. Western blot analysis of TLR1 using anti-TLR1 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat spleen tissue lysates, Lane 2: mouse small intestine tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TLR1 antigen affinity purified polyclonal antibody (Catalog # ) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TLR1 at approximately 90KD. The expected band size for TLR1 is at 90KD.





#### **Immunohistochemistry**

Image 2. IHC analysis of TLR1 using anti-TLR1 antibody . TLR1 was detected in paraffin-embedded section of mouse spleen tissue . Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2μg/ml rabbit anti-TLR1 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

#### **Immunohistochemistry**

**Image 3.** IHC analysis of TLR1 using anti-TLR1 antibody . TLR1 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2μg/ml rabbit anti-TLR1 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog #SA1022) with DAB as the chromogen.

Please check the product details page for more images. Overall 4 images are available for ABIN5693146.