

Datasheet for ABIN5693294
anti-WWOX antibody (AA 1-245)



[Go to Product page](#)

5 Images

Overview

Quantity:	100 µg
Target:	WWOX
Binding Specificity:	AA 1-245
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WWOX antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Brand:	Picoband™
Immunogen:	E. coli-derived human WWOX recombinant protein (Position: M1-D245).
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	Rabbit IgG polyclonal antibody for WWOX detection. Tested with WB, IHC-P, Direct ELISA in Human, Mouse, Rat.

Target Details

Target:	WWOX
Alternative Name:	WWOX (WWOX Products)
Background:	Synonyms: WW domain-containing oxidoreductase, Fragile site FRA16D oxidoreductase, Short

Target Details

chain dehydrogenase/reductase family 41C member 1, WWOX, FOR, SDR41C1, WOX1

Tissue Specificity: Widely expressed. Strongly expressed in testis, prostate, and ovary.

Overexpressed in cancer cell lines. Isoform 5 and isoform 6 may only be expressed in tumor cell lines.

Background: WW domain-containing oxidoreductase is an enzyme that in humans is encoded by the WWOX gene. This gene encodes a member of the short-chain dehydrogenases/reductases (SDR) protein family. It spans the FRA16D common chromosomal fragile site and appears to function as a tumor suppressor gene. Expression of the encoded protein is able to induce apoptosis, while defects in this gene are associated with multiple types of cancer. Disruption of this gene is also associated with autosomal recessive spinocerebellar ataxia 12. Disruption of a similar gene in mouse results in impaired steroidogenesis, additionally suggesting a metabolic function for the protein. Alternative splicing results in multiple transcript variants.

Application Details


Application Notes:	Recommended Detection Systems: Enhanced Chemiluminescent Kit with anti-Rabbit IgG (ABIN921124) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
	Application Details: Western blot, 0.1-0.5 µg/mL
	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/mL
	Direct ELISA, 0.1-0.5 µg/mL
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , 0.05 mg NaN ₃ .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month.


It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Images




Western Blotting

Image 1. Western blot analysis of WWOX using anti-WWOX antibody . Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: human MCF-7 whole cell lysates, Lane 2: rat brain tissue lysates, Lane 3: rat kidney tissue lysates, Lane 4: rat testis tissue lysates, Lane 5: mouse brain tissue lysates, Lane 6: mouse kidney tissue lysates, Lane 7: mouse testis tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-WWOX antigen affinity purified polyclonal antibody (Catalog #) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for WWOX at approximately 46KD. The expected band size for WWOX is at 46KD.



Immunohistochemistry

Image 2. IHC analysis of WWOX using anti-WWOX antibody . WWOX was detected in paraffin-embedded section of human prostatic cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1µg/ml rabbit anti-WWOX Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



Immunohistochemistry

Image 3. IHC analysis of WWOX using anti-WWOX antibody . WWOX was detected in paraffin-embedded section of mouse brain tissue . Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1µg/ml rabbit anti-WWOX Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN5693294.