

Datasheet for ABIN5706396

Universal Blocking Buffer for Western Blotting



[Go to Product page](#)

1 Image

Overview

Quantity:	500 mL
Application:	Western Blotting (WB)

Product Details

Purification:	Blocking buffer was prepared using ultra pure reagents dissolved in pharmaceutical grade water (WFI) and consists of a proprietary protein formulation in TRIS buffered saline at pH 7.6 with thimerisol added as an antimicrobial agent.
Sterility:	Aseptic filtered

Application Details

Application Notes:	<p>Application Note: It allows for superior signal detection and lower background noise in colorimetric, chemiluminescent, and fluorescent western blotting and other applications making it a convenient general purpose blocking agent. Antibody conjugates prepared with IRDye® 800 and IRDye® 700DX (Licor), Cy2™, Cy3™, Cy3.5™, Cy5™ and Cy5.5™ (GE Healthcare), DyLight™ 405, DyLight™ 549, DyLight™ 649, DyLight™ 680, and DyLight™ 800 (Thermo Fisher/Pierce) and Alexa Fluor® 488, Alexa Fluor® 532, Alexa Fluor® 546, Alexa Fluor® 647 and Alexa Fluor® 680 (Invitrogen/Molecular Probes) and ATTO (Atto-Tec) have been validated on various platforms using this product with superior results compared to other commercially available products. In the infrared range, where little to no autofluorescence occurs, specific signal is sharply evident from any background giving the best possible signal-to-noise ratio. This allows for detection levels in the picogram range which rivals the sensitivity of chemiluminescence on film for western blotting. Superior results are also seen when this product is used for simultaneous labeling (multiplex) in western blots or microscopy using various fluorochrome combinations for multicolor imaging. Membranes blocked with this</p>
--------------------	---

Application Details

Blockig Buffer can be dried and are very stable.

Western Blot Dilution: User Defined

Comment:

Synonyms: Multiplex Blocking Buffer, Immunoblot Blocking Buffer, Blocking Solution, Blocking Buffer Western Blot, Western Blot Blocking Buffer, Alexa Dye Blocking Buffer, DyLight Blocking Buffer, colorimetric, chemiluminescent blocking buffer, WB block

Background: It is an universal blocking buffer solution for Western Blotting. It is specifically designed for colorimetric, chemiluminescent, and fluorescent western blotting. Universal Blocking Buffer is recommended for blocking when phospho specific antibodies are used. Pure nitrocellulose membrane is recommended for maximum performance. Other membranes, such as PVDF or nitrocellulose embedded in a support can be used, but may generate elevated backgrounds. Protein should be transferred from gel to membrane using standard protocols. Blocking buffer can be used for membrane blocking and to dilute both primary and secondary antibodies. Western Blot blocking buffer is suitable for use with chemiluminiscent and fluorescent western blot imaging systems produced by Bio-Rad Laboratories, GE Healthcare, Alpha Innotech, FujiFilm Life Science, Licor Biosciences, UVP and Syngene.

Restrictions:

For Research Use only

Handling

Format:

Liquid

Concentration:

1 X

Preservative:

Thimerosal (Merthiolate)

Precaution of Use:


This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

4 °C

Expiry Date:

6 months



Western Blotting

Image 1. Effect of selected cholinergic agonists, anthelmintics and antagonists on the Asu-EAT-2 receptor expressed in *Xenopus* oocytes. (D) Localization of Asu-eat-2 and Asu-eat-18 mRNA in different body tissues of the *A. suum* worm (n = 5). RT-PCR analysis of Asu-eat-2 (lanes 2, 5, 8, 11) and Asu-eat-18 (lanes 3, 6, 9, 12) and gapdh control (lanes 4, 7, 10, 13) in pharynx, ovijector, head, and gut region. The PCR product sizes for eat-2, eat-18 and gapdh were 949, 213 and 411 bp respectively. Lane 1, FastRuler High Range DNA ladder. The gels were blotted onto PVDF membranes and blocked with BlockOut blocking buffer (p/n MB-073). Fig 3. PMID: 32243475.