

Datasheet for ABIN5706798

## anti-Histone H4 antibody (2meLys20)



[Go to Product page](#)

### 5 Images

#### Overview

Quantity:	50 µg
Target:	Histone H4
Binding Specificity:	2meLys20
Reactivity:	Human, C. elegans
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Histone H4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Chromatin Immunoprecipitation (ChIP), Dot Blot (DB), Fluorescence Microscopy (FM)

#### Product Details

Purpose:	Histone H4 K20me2 Antibody
Immunogen:	Immunogen: Histone H4 [Dimethyl Lys20] affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic dimethylated peptide surrounding Lysine 20 of human Histone H4. Immunogen Type: Conjugated Peptide
Isotype:	IgG
Cross-Reactivity (Details):	This antibody reacts with human Histone H4.
Characteristics:	Synonyms: rabbit anti-Histone H4 dimethyl Lys20 antibody, H4K20me2, HIST2H4B, HIST1H4H, HIST1H4I, HIST1H4J, HIST1H4K, HIST1H4L, HIST2H4, HIST2H4A, HIST1H4A, HIST1H4B, HIST1H4C, HIST1H4D, HIST1H4E, HIST1H4F, histone cluster 4, H4, histone 4, H4

## Product Details

Purification:	Anti-Histone H4 [Dimethyl Lys20] was affinity purified from monospecific antiserum by immunoaffinity chromatography.
---------------	--

Sterility:	Sterile filtered
------------	------------------

## Target Details

Target:	Histone H4
---------	------------

Abstract:	<a href="#">Histone H4 Products</a>
-----------	-------------------------------------

Background:	<p>Background: When K20 is di-methylated on H4, the downstream effect is silencing of genes. This modification is also necessary and sufficient for 53BP1 binding, which is a prerequisite for DNA repair, a highly conserved mechanism. H4K20me2 is associated with hypoacetylation, and inactivation of certain genes. NSD1 and Suv420h1/2 trigger the dimethylation of K20, but only after SET8 catalyzes the mono-methylation of the same lysine, indicating a complex regulation of this modification that can so seriously affect replication and stability of genomic information. Anti-Histone H4 are ideal for researchers interested in Chromatin Modifiers, Chromatin Research, DNA Repair, DNA replication Transcription Translation and Splicing, Histones and Modified Histones, and Epigenetics research.</p>
-------------	--

Gene ID:	121504
----------	--------

NCBI Accession:	<a href="#">NP_001029249</a>
-----------------	------------------------------

UniProt:	<a href="#">P62805</a>
----------	------------------------

## Application Details

Application Notes:	<p>Immunohistochemistry Dilution: 1:50</p> <p>Application Note: Anti-Histone H4 [Dimethyl Lys20] antibody is tested for Western Blot, Dot Blot, Chromatin Immunoprecipitation, Immunofluorescence, and Immunocytochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~13 kDa corresponding to Histone H4 protein by Western Blotting in HeLa histone prep lysate or the appropriate cell lysate or extract. Epi-Plus™ antibody production in collaboration with Novus Biologicals.</p> <p>ChIP Dilution: 2-5 µg/million cells</p> <p>Western Blot Dilution: 1:2000</p> <p>IF Microscopy Dilution: 1:50</p> <p>Other: Dot Blot 1:1000</p>
--------------------	---


## Application Details

Restrictions: For Research Use only

## Handling


Format:	Liquid
Concentration:	0.53 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 30 % Glycerol Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months

## Images




### Dot Blot

**Image 1.** Dot Blot of Rabbit Histone H4 [Dimethyl Lys20] Antibody. Lane 1: K9-K14. Lane 2: K9-Kac-K14-Kac. Lane 3: K9-K14-Kac. Lane 4: K9-ac-K14. Load: 0.3, 1, 3, 10, and 30 picomoles of peptide. Primary antibody: Histone H4 [Dimethyl Lys20] antibody at 1:2000 for 45 min at 4 °C. Secondary antibody: Dylight™488 rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5 % BLOTTO overnight at 4 °C.



### Fluorescence Microscopy

**Image 2.** Immunofluorescence of Rabbit Anti-Histone H4 [Dimethyl Lys20] Antibody. Tissue: HeLa cells. Fixation: 0.5 % PFA. Antigen retrieval: Not required. Primary antibody: Histone H4 [Dimethyl Lys20] antibody at a 1:50 dilution for 1 h at RT. Secondary antibody: FITC secondary antibody at 1:10,000 for 45 min at RT. Localization: Histone H4 [Dimethyl Lys20] is nuclear and chromosomal. Staining: Histone H4 [Dimethyl Lys20] is expressed in green, nuclei are counterstained with DAPI (blue).



### Western Blotting

**Image 3.** Western Blot of Rabbit Anti-Histone H4 [Dimethyl Lys20] Antibody. Lane 1: C. elegans embryo lysate. Load: 30 µg per lane. Primary antibody: Histone H4 [Dimethyl Lys20] at 1:2000 for overnight at 4 °C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5 % BLOTTO overnight at 4 °C. Predicted/Observed size: ~13 kDa. Other band(s): None.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN5706798.