

## Datasheet for ABIN5707409

# Goat anti-Rabbit IgG (Fc Region) Antibody (Biotin)





Go to Product page

U	V	e	rV	16	9	Λ

Quantity:	0.5 mg
Target:	IgG
Binding Specificity:	Fc Region
Reactivity:	Rabbit
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, FLISA
Product Details	
Product Details  Isotype:	IgG
	IgG  Reacts with the Fc region of rabbit IgG
Isotype:	<u> </u>
Isotype: Specificity:	Reacts with the Fc region of rabbit IgG
Isotype: Specificity: Characteristics:	Reacts with the Fc region of rabbit IgG  Goat Anti-Rabbit IgG Fc-BIOT
Isotype: Specificity: Characteristics:	Reacts with the Fc region of rabbit IgG  Goat Anti-Rabbit IgG Fc-BIOT  Purification Method: Affinity chromatography on pooled rabbit IgG covalently linked to
Isotype: Specificity: Characteristics: Purification:	Reacts with the Fc region of rabbit IgG  Goat Anti-Rabbit IgG Fc-BIOT  Purification Method: Affinity chromatography on pooled rabbit IgG covalently linked to
Isotype: Specificity: Characteristics: Purification:  Target Details	Reacts with the Fc region of rabbit IgG  Goat Anti-Rabbit IgG Fc-BIOT  Purification Method: Affinity chromatography on pooled rabbit IgG covalently linked to agarose.

### **Application Details**

Ann	lication	Notes:
$\neg$ pp	iication	INOLUS.

- · Applications: Quality tested applications include ELISA FLISA
- Other referenced applications include WB
  - **Working Dilutions:** ELISA Purified (UNLB) antibody 1 g/mL HRP conjugate 1:2,000 1:4,000 BIOT conjugate 1:5,000 1:20,000 FLISA FITC conjugate 1:200 1:400


Restrictions:

For Research Use only

#### Handling

Concentration:	0.5 mg/mL
Buffer:	0.5 mg in 1.0 mL of PBS/Sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 2-8°C

#### **Images**



#### **ELISA**

**Image 1.** ELISA plate was coated with purified rabbit IgG Fc and IgG Fab. Immunoglobulins were detected with serially diluted Goat Anti-Rabbit IgG Fc-BIOT.