# antibodies -online.com





# anti-TAF7L antibody (Internal Region)



Image



Go to Product page

$\sim$						
	1//	Д	r۱	1	<b>Θ</b> 1	٨

Quantity:	100 μg	
Target:	TAF7L	
Binding Specificity:	Internal Region	
Reactivity:	Human	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	This TAF7L antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	

#### **Product Details**

Purpose:	TAF7L			
Immunogen:	Peptide with sequence C-NHFQSVLEQLELQEK, from the internal region (near C Terminus) of the protein sequence according to NP_079161.2.			
Sequence:	NHFQSVLEQL ELQEK			
Isotype:	IgG			
Cross-Reactivity:	Human			
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.			
Grade:	Verified			

## **Target Details**

rarget Details			
Target:	TAF7L		
Alternative Name:	TAF7L (TAF7L Products)		
Background:	TAF7L, TAF7-like RNA polymerase II, TATA box binding protein (TBP)-associated factor, 50 kDa FLJ23157, TAF2Q, dJ738A13.1, OTTHUMP00000023675, TAF7-like RNA polymerase II, TATA box binding protein (TBP)-associated factor, 50 kD, TATA box binding protein (T		
Gene ID:	54457		
NCBI Accession:	NP_079161		
Application Details			
Application Notes:	Western Blot: Approx 40 kDa band observed in Human Testis lysates (calculated MW of 52.6 kDa according to NP_079161.2). This molecular weight is routinely observed by other sources. Recommended concentration: 0.3-1 µg/mL.  Peptide ELISA: antibody detection limit dilution 1:16000.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	0.5 mg/mL		
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Handling Advice:	Minimize freezing and thawing.		
Storage:	-20 °C		
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.		

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

### **Western Blotting**

**Image 1.** ABIN570772 (1 $\mu$ g/ml) staining of Human Testis lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.