antibodies .- online.com







anti-TNFSF15 antibody





Go to Product page

\sim	
()\/\	view
	V I C V V

Alternative Name:

Background:

Quantity:	100 μg
Target:	TNFSF15
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This TNFSF15 antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	A full length human recombinant protein was used as the immunogen for this recombinant TNFSF15 antibody.
Clone:	
	VEGI-2052R
Isotype:	VEGI-2052R IgG kappa
Isotype:	IgG kappa
Isotype: Purification:	IgG kappa Purified

TNF ligand-related molecule 1 is an anti-angiogenic cytokine that is also called Tumor necrosis

TNFSF15 / VEGI / TL1A (TNFSF15 Products)

factor superfamily, member 15 (TNFSF15) and Vascular endothelial growth inhibitor (VEGI/VEGI192A). This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21/DR6. It can activate NF-kappaB and MAP kinases, and acts as an autocrine factor to induce apoptosis in endothelial cells. This cytokine is also found to inhibit endothelial cell proliferation, and thus may function as an angiogenesis inhibitor. Reduced expression of TL1A/TNFSF15/VEGI has been reported as a marker of poor prognosis in breast cancer.

Gene ID:

9966

Pathways:

Positive Regulation of Endopeptidase Activity, Autophagy

Application Details

Application Notes:

Titration of the recombinant TNFSF15 antibody may be required for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.\. Immunohistochemistry (FFPE): 1-2 μ g/mL,Prediluted IHC only format : incubate for 30 min at RT (1)


Restrictions:

For Research Use only

Handling

Buffer:	1 mg/mL in 1X PBS, BSA free, sodium azide free
Preservative:	Azide free
Storage:	4 °C,-20 °C
Storage Comment:	Store the recombinant TNFSF15 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).


Human Protein Microarray Specificity Validation



Microarray


Image 1. Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using recombinant TNFSF15 antibody (clone VEGI/2052R). These results demonstrate the foremost specificity of the VEGI/2052R mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



Immunohistochemistry

Image 2. IHC testing of FFPE human spleen with recombinant TNFSF15 antibody (clone VEGI/2052R). Required HIER: boil sections in 10mM Tris with 1mM EDTA, pH9, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE

Image 3. SDS-PAGE analysis of purified, BSA-free recombinant TNFSF15 antibody (clone VEGI/2052R) as confirmation of integrity and purity.

Please check the product details page for more images. Overall 4 images are available for ABIN5708012.