antibodies .- online.com







anti-FANCE antibody (Internal Region)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μg
Target:	FANCE
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This FANCE antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Purpose:	FANCE
Immunogen:	C-EHKSLESLADGGS
Sequence:	EHKSLESLAD GGS
Isotype:	IgG
Cross-Reactivity:	Cow, Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

Target:	FANCE
Alternative Name:	FANCE (FANCE Products)
Background:	FANCE, Fanconi anemia, complementation group E, FACE, FAE
Gene ID:	2178
NCBI Accession:	NP_068741
Pathways:	DNA Damage Repair

Application Details

		Peptide ELISA: antibody detection limit dilution 1:32000.
		explanation in the literature for the band we observe given the calculated
		Uterus lysates after 0.1 μ g/mL antibody staining. Please note that currently we cannot find an
		Western Blot: Preliminary experiments gave an approx 70 kDa band in Human Adipose and
		trophoblasts. Recommended concentration, 3-6 µg/mL.
Application Notes: Immunohistochemistry: In paraffin embedded Human Place		Immunohistochemistry: In paraffin embedded Human Placenta shows nuclear staining in some

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.