

Datasheet for ABIN570874  
**anti-PTPRB antibody (Internal Region)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µg
Target:	PTPRB
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This PTPRB antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Flow Cytometry (FACS)

## Product Details

Purpose:	PTPRB
Immunogen:	C-ENHSFSQERTVPDK
Sequence:	ENHSFSQERT VPK
Isotype:	IgG
Cross-Reactivity:	Cow, Dog, Human, Pig, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

## Target Details



Target:	PTPRB
Alternative Name:	PTPRB ( <a href="#">PTPRB Products</a> )
Background:	PTPRB, protein tyrosine phosphatase, receptor type, B, DKFZp686E2262, DKFZp686H15164, FLJ44133, HPTP-BETA, HPTPB, MGC142023, MGC59935, PTPB, R-PTP-BETA, VEPTP, protein tyrosine phosphatase, receptor type, beta polypeptide
Gene ID:	5787, 314843
NCBI Accession:	<a href="#">NP_001103224</a> , <a href="#">NP_002828</a> , <a href="#">NP_001193901</a>

## Application Details

Application Notes:	Peptide ELISA: antibody detection limit dilution 1:16000.
Comment:	<b>Immunofluorescence:</b> Strong expression of the protein seen in the membranes and vesicles of A431 cells. Recommended concentration: 10µg/ml. <b>Flow Cytometry:</b> Flow cytometric analysis of A431 cells. Recommended concentrati
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



### Immunofluorescence

**Image 1.** ABIN570874 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

### Flow Cytometry

**Image 2.** ABIN570874 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.