

Datasheet for ABIN571066
anti-GPR17 antibody (C-Term)

6 Images

[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	GPR17
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This GPR17 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)

Product Details

Purpose:	GPR17
Immunogen:	Peptide with sequence C-EGKTNESSLSAKSE, from the C Terminus of the protein sequence according to NP_005282.1.
Sequence:	EGKTNESSLS AKSE
Isotype:	IgG
Cross-Reactivity:	Cow, Human, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details


Target:	GPR17
Alternative Name:	GPR17 (GPR17 Products)
Background:	GPR17, G protein-coupled receptor 17, DKFZp686M18273
Gene ID:	2840, 574402, 767613
NCBI Accession:	NP_005282

Application Details

Application Notes:	Western Blot: Approx 40 kDa band observed in Mouse Brain lysates (calculated MW of 37.8 kDa according to Mouse NP_001020552.1). Recommended concentration: 2-3 µg/mL. Primary incubation 1 hour at room temperature. Peptide ELISA: antibody detection limit dilution 1:4000.
Comment:	Immunofluorescence: Strong expression of the protein seen in the vesicles of HeLa cells and additionally in the plasma membrane of U251 cells. Recommended concentration: 10µg/ml.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.




Immunofluorescence

Image 1. ABIN571066 Immunofluorescence analysis of paraformaldehyde fixed U251 cells, permeabilized with 0.15 % Triton. Primary incubation 1hr (10 µg/mL) followed by Alexa Fluor 488 secondary antibody (2 µg/mL), showing membrane and vesicle staining. The nuclear stain i



Western Blotting

Image 2. ABIN571066 (0.1µg/ml) staining of Human Cerebellum lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



Western Blotting

Image 3. Antibody (0.3µg/ml) staining of Rat Brain lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN571066.