antibodies -online.com





anti-STAG2 antibody (Internal Region)

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	STAG2
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This STAG2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	Stromal antigen 2 / STAG2
Immunogen:	Peptide with sequence C-SRGSTVRSKKSKPST, from the internal region of the protein sequence according to NP_001036214.1, NP_006594.3.
Sequence:	SRGSTVRSKK SKPST
Isotype:	IgG
Specificity:	This antibody is expected to recognize both reported isoforms (NP_001036214.1, NP_006594.3). Reported variants NP_001036215.1 and NP_001036214.1 represent identical protein: Reported variants NP_001036216.1 and NP_006594.3 represent identical protein.
Cross-Reactivity:	Cow, Dog, Human, Mouse, Pig, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

Product Details chromatography using the immunizing peptide. Grade: Verified Target Details STAG2 Target: Alternative Name: STAG2 (STAG2 Products) Background: STAG2, stromal antigen 2, DKFZp686P168, DKFZp781H1753, FLJ25871, SA-2, SA2, bA51701.1, OTTHUMP00000024338, OTTHUMP00000043514, OTTHUMP00000043515, OTTHUMP00000081754, OTTHUMP00000081755, SCC3 homolog 2 Gene ID: 10735, 20843, 313304 NCBI Accession: NP_001036214, NP_006594 Stem Cell Maintenance Pathways: **Application Details** Western Blot: Approx 160 kDa band observed in lysates of cell line K562 and ~140 kDa in **Application Notes:** rodent Spleen while ~150 kDa in Mouse Thymus lysates (calculated MW of 146 kDa according to Human NP_001036214.1 and 140 kDa according to Mouse NP_001071180.1). Recommen Peptide ELISA: antibody detection limit dilution 1:32000. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin. Preservative: Sodium azide

should be handled by trained staff only.

Minimize freezing and thawing.

-20 °C

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Precaution of Use:


Handling Advice:

Storage:

Storage Comment:

Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.

Images





Western Blotting

Image 1. ABIN571073 (0.3μg/ml) staining of Mouse Spleen (A), Mouse Thymus (B) and Rat Spleen (C) lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Western Blotting

Image 2. ABIN571073 (0.03μg/ml) staining of K562 lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.