antibodies - online.com







anti-MECOM antibody (Internal Region)



Image



Overview

Quantity:	100 μg
Target:	MECOM
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This MECOM antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	EVI1 / AML1
Immunogen:	Peptide with sequence C-DKESLHSTSH, from the internal region of the protein sequence according to NP_001098547.3, NP_005232.2, NP_004982.2, NP_001157471.1, NP_001157472.1.
Sequence:	DKESLHSTSH
Isotype:	IgG
Specificity:	This antibody is expected to recognize all reported isoforms (NP_001098547.3, NP_005232.2, NP_004982.2, NP_001157471.1, NP_001157472.1). Reported variants represent identical protein: NP_001098548.2, NP_005232.2
Cross-Reactivity:	Human

Product Details	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity
	chromatography using the immunizing peptide.
Grade:	Verified
Target Details	
Target:	MECOM
Alternative Name:	MECOM (MECOM Products)
Background:	MECOM, MDS1 and EVI1 complex locus, AML1-EVI-1, EVI1, MDS1, MDS1-EVI1, MGC163392, MGC97004, PRDM3, AML1-EVI-1 fusion protein, ecotropic viral integration site 1, myelodysplasia syndrome 1, myelodysplasia syndrome-associated protein 1, oncogene EVI1, zinc
Gene ID:	2122, 14013
NCBI Accession:	NP_001098547, NP_005232, NP_004982, NP_001157471, NP_001157472
Application Details	
Application Notes:	Western Blot: Approx 170 kDa band observed in lysates of cell lines A549 and HepG2
	(calculated MW of 126 kDa according to NP_001098547.3). This molecular weight is routinely
	observed by other sources. Recommended concentration: 1-3 μ g/mL.
	Peptide ELISA: antibody detection limit dilution 1:16000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.

Handling

Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.
1	

Images

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Western Blotting

Image 1. ABIN571139 (2μg/ml) staining of A549 lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.