antibodies -online.com





anti-TRIP6 antibody (Internal Region)



Image



Go to Product page

Overview

Quantity:	100 μg
Target:	TRIP6
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This TRIP6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	TRIP6
Immunogen:	C-SHGVLQHTQGLP
Sequence:	SHGVLQHTQG LP
Isotype:	IgG
Cross-Reactivity:	Cow, Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

rarget Details	
Target:	TRIP6
Alternative Name:	TRIP6 (TRIP6 Products)
Background:	TRIP6, thyroid hormone receptor interactor 6, MGC10556, MGC10558, MGC29959, MGC3837
	MGC4423, OIP1, ZRP-1, OPA-interacting protein 1, thyroid receptor-interacting protein 6, zyxir
	related protein 1
Gene ID:	7205
NCBI Accession:	NP_003293
Pathways:	Cell-Cell Junction Organization
Application Details	
Application Notes:	Western Blot: Approx 50 kDa band observed in lysates of cell line K562 (calculated MW of
	50.3 kDa according to NP_003293.2). Recommended concentration: 1-3 µg/mL.
	Peptide ELISA: antibody detection limit dilution 1:16000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum
	albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerate
	at 4°C for a few weeks and still remain viable.



Western Blotting

Image 1. ABIN571141 ($2\mu g/ml$) staining of K562 lysate (35 μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.