antibodies -online.com





anti-HCN3 antibody (Internal Region)





Go to Product page

()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μg	
Target:	HCN3	
Binding Specificity:	Internal Region	
Reactivity:	Rat	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	This HCN3 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	

Product Details

Purpose:	HCN3	
Immunogen:	C-RGVRGRAPSTGAQ	
Sequence:	RGVRGRAPST GAQ	
Isotype:	IgG	
Cross-Reactivity:	Cow, Dog, Human, Mouse, Rat	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.	
Grade:	Verified	

Target Details

rarget Details		
Target:	HCN3	
Alternative Name:	HCN3 (HCN3 Products)	
Background:	HCN3, hyperpolarization activated cyclic nucleotide-gated potassium channel 3, KIAA1535, MGC131493, potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 3	
Gene ID:	57657, 15168	
NCBI Accession:	NP_065948	
Application Details		
Application Notes:	Western Blot: Approx 85 kDa band observed in Rat Kidney lysates (calculated MW of 86.0 kDa according to Human NP_065948.1 and 86.8 kDa according to Rat NP_446137.1). Recommended concentration: 1-3 µg/mL. Peptide ELISA: antibody detection limit dilution 1:32000.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Minimize freezing and thawing.	
Storage:	-20 °C	
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.	

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Western Blotting

Image 1. ABIN571196 (1 μ g/ml) staining of Rat Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.