antibodies

Datasheet for ABIN5712073 ATPC1 Protein (AA 43-161, partial) (GST tag)



Overview

Image

Quantity:	100 µg
Target:	ATPC1
Protein Characteristics:	AA 43-161, partial
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This ATPC1 protein is labelled with GST tag.
Application:	SDS-PAGE (SDS)

Product Details

Sequence:	ASPTQVFFNG ANVRQVDVPT LTGAFGILAA HVPTLQVLRP GLVVVHAEDG TTSKYFVSSG SIAVNADSSV QLLAEEAVTL DMLDLGAAKA NLEKAQAELV GTADEATRAE IQIRIEANE
Purification:	SDS-PAGE
Purity:	> 90 %

Target Details

Target:	ATPC1
Alternative Name:	ATPD (ATPC1 Products)
Background:	Mitochondrial mbrane ATP synthase (F1F0 ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the mbrane which is generated by electron
	transport complexes of the respiratory chain. F-type ATPases consist of two structural

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5712073 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

	domains, F1 - containing the extrambraneous catalytic core, and F0 - containing the mbrane
	proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP
	turnover in the catalytic domain of F1 is coupled via a rotary mechanism of the central stalk
	subunits to proton translocation. Part of the complex F1 domain and of the central stalk which
	is part of the complex rotary elent. Rotation of the central stalk against the surrounding
	alpha3beta3 subunits leads to hydrolysis of ATP in three separate catalytic sites on the beta
	subunits.
Molecular Weight:	39.8 kDa
UniProt:	P30049
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.1-2 mg/mL
Buffer:	20 mM Tris-HCl based buffer, pH 8.0
Storage:	-80 °C,4 °C,-20 °C
Storage Comment:	Store at -20°C, for extended storage, conserve at -20°C or -80°C. Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.




Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN5712073 | 09/10/2023 | Copyright antibodies-online. All rights reserved.