

Datasheet for ABIN571350

anti-KIT antibody



[Go to Product page](#)

1 Image

7 Publications

Overview

Quantity:	100 µg
Target:	KIT
Reactivity:	Human, Cow, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This KIT antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Purpose:	Anti-Hu CD117 Purified
Immunogen:	MOLM-1 megakaryocytic cells
Clone:	104D2
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 104D2 detects extracellular part of CD117 / c-Kit protooncogen.
Cross-Reactivity (Details):	Human, Non-Human Primates, Bovine
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	KIT
Alternative Name:	CD117 (KIT Products)
Background:	KIT proto-oncogene, receptor tyrosine kinase,CD117 / c-Kit (stem cell factor receptor) is a 145 kDa receptor tyrosine kinase that regulates cell proliferation, adhesion, chemotaxis, apoptosis and other cell processes. Mutations of CD117 / c-Kit can lead to growth and progression of tumours. After binding of its ligand, SCF (stem cell factor), CD117 / c-Kit is autophosphorylated on its intracellular domains and activated. CD117 is expressed on pluripotent hematopoietic progenitor cells, mast cells and various cancer cells, e.g. acute myeloid leukemia cells.,c-Kit, PBT, SCFR
Gene ID:	3815
UniProt:	P10721
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Sensory Perception of Sound , Stem Cell Maintenance , Production of Molecular Mediator of Immune Response , Regulation of long-term Neuronal Synaptic Plasticity

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1 µg/mL.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Product cited in: Stevenson, McGlynn, Hodge, McLinden, George, Davies, Shiels: "Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations." in: **Stem cells and development**, Vol. 18, Issue 10, pp. 1389-98, (2009) ([PubMed](#)).

Wihlidal, Varga, Pfeilstöcker, Karlic: "Expression and functional significance of osteocalcin splicing in disease progression of hematological malignancies." in: **Leukemia research**, Vol. 30, Issue 10, pp. 1241-8, (2006) ([PubMed](#)).


Blair, Sutherland: "Primitive acute myeloid leukemia cells with long-term proliferative ability in vitro and in vivo lack surface expression of c-kit (CD117)." in: **Experimental hematology**, Vol. 28, Issue 6, pp. 660-71, (2000) ([PubMed](#)).

Yoshino, Ami, Terao, Tashiro, Honda: "Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies." in: **Experimental animals / Japanese Association for Laboratory Animal Science**, Vol. 49, Issue 2, pp. 97-110, (2000) ([PubMed](#)).

Broudy, Lin, Liles, Corey, OLaughlin, Mou, Linnekin: "Signaling via Src family kinases is required for normal internalization of the receptor c-Kit." in: **Blood**, Vol. 94, Issue 6, pp. 1979-86, (1999) ([PubMed](#)).

There are more publications referencing this product on: [Product page](#)

Images



Flow Cytometry

Image 1. Surface staining of human peripheral blood cells with anti-CD117 (104D2) purified.