



[Go to Product page](#)

Datasheet for ABIN574352

## anti-TFF3 antibody (AA 15-74)

### 3 Images

#### Overview

Quantity:	100 µg
Target:	TFF3
Binding Specificity:	AA 15-74
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### Product Details

Brand:	IHC-plus™
Immunogen:	TFF3 (AAH17859, 15 a.a. ~ 74 a.a) full length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.  Type of Immunogen: Recombinant protein
Clone:	3D9
Isotype:	IgG1 kappa
Purification:	Purified from ascites by Protein A

#### Target Details

Target:	TFF3
---------	------

## Target Details

---

Alternative Name: TFF3 / Trefoil Factor 3 ([TFF3 Products](#))

---

Background: Name/Gene ID: TFF3

Synonyms: TFF3, HP1.B, Intestinal trefoil factor, HITF, ITF, TFI, p1B, Polypeptide P1.B, Trefoil factor 3, Trefoil factor 3 (intestinal)

---

Gene ID: 7033

## Application Details

---

Application Notes: Approved: ELISA, IHC, IHC-P (5 µg/mL), WB

Usage: Western blot performed with the recombinant protein used as the immunogen.  
Sandwich ELISA (Recombinant protein).

---

Comment: Target Species of Antibody: Human

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: Lot specific

---

Buffer: PBS, pH 7.4

---


Handling Advice: Aliquot to Avoid freeze/thaw cycles.

---

Storage: -20 °C


---

Storage Comment: Store at -20°C. Aliquot to avoid freeze/thaw cycles.




### Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)

**Image 1.** Human Small Intestine: Formalin-Fixed, Paraffin-Embedded (FFPE)



### Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)

**Image 2.** Human Uterus: Formalin-Fixed, Paraffin-Embedded (FFPE)



### Immunohistochemistry

**Image 3.** Anti-TFF3 antibody IHC of human small intestine. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.