

Datasheet for ABIN5854584

## CD80 Protein (CD80) (AA 37-246) (His tag)



[Go to Product page](#)

### 1 Image

#### Overview

|                               |   |
|-------------------------------|---|
| Quantity:                     | 50 µg                                       |
| Target:                       | CD80  |
| Protein Characteristics:      | AA 37-246                                   |
| Origin:                       | Mouse                                       |
| Source:                       | Baculovirus infected Insect Cells           |
| Protein Type:                 | Recombinant                                 |
| Biological Activity:          | Active                                      |
| Purification tag / Conjugate: | This CD80 protein is labelled with His tag. |
| Application:                  | SDS-PAGE (SDS), Activity Assay (AcA)        |

#### Product Details

|                              |  |
|------------------------------|--|
| Sequence:                    | DVDEQLSKSV KDKVLLPCRY NSPHEDESED RIYWQKHDKV VLSVIAGKLK VWPEYKNRTL<br>YDNTTYSLLI LGLVLSDRGT YSCVVQKKER GTYEVKHLAL VKLSIKADFS TPNITESGNP<br>SADTKRITCF ASGGFPPKPRF SWLENGRELP GINTTISQDP ESELYTISSQ LDFNTTRNHT<br>IKCLIKYGDA HVSEDFTWEK PPEDPPDSKN |
| Purity:                      | > 95% by SDS-PAGE  |
| Endotoxin Level:             | < 1 EU per 1 µg of protein (determined by LAL method)  |
| Biological Activity Comment: | Measured by its binding ability in a functional ELISA with Mouse CTLA4 (CAT# ATGP4046).  |

#### Target Details

|         |      |
|---------|------|
| Target: | CD80 |
|---------|------|

## Target Details


|                   |  |
|-------------------|--|
| Alternative Name: | B7-1/CD80 ( <a href="#">CD80 Products</a> )  |
| Background:       | <p>Cd80, also known as T-lymphocyte activation antigen CD80, is a member of cell surface immunoglobulin superfamily and is expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. It is the ligand for two different proteins on the T cell surface: Cd28 (for autoregulation and intercellular association) and Ctla4 (for attenuation of regulation and cellular disassociation). This protein also plays a role in induction of innate immune responses by activating NF-KB-signaling pathway in macrophages. It is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas. Recombinant mouse Cd80, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.</p> |
| Molecular Weight: | 24.7 kDa (216aa)   |
| NCBI Accession:   | <a href="#">NP_033985</a>  |
| UniProt:          | <a href="#">Q00609</a>   |
| Pathways:         | <a href="#">TCR Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">EGFR Signaling Pathway</a> , <a href="#">Neurotrophin Signaling Pathway</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Cancer Immune Checkpoints</a>   |

## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions:      | For Research Use only  |

## Handling

|                  |   |
|------------------|---|
| Format:          | Liquid  |
| Concentration:   | 0.25 mg/mL  |
| Buffer:          | Liquid. In Phosphate Buffered Saline ( pH 7.4) containing 10 % glycerol.  |
| Storage:         | 4 °C,-20 °C,-80 °C  |
| Storage Comment: | Can be stored at +2°C to +8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Avoid repeated freezing and thawing cycles. |



| SDS-PAGE |
|----------|
| Image 1. |