

Datasheet for ABIN605079
anti-OTUD4 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	0.05 mg
Target:	OTUD4
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OTUD4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Isotype:	IgG
Specificity:	Multiple isoforms of OTUD4 are known to exist. OTUD4 antibody is predicted to not cross-react with other members of the OTUD family.
Purification:	Immunoaffinity purified

Target Details

Target:	OTUD4
Alternative Name:	OTUD4 (OTUD4 Products)

Target Details


Background:	Name/Gene ID: OTUD4 Synonyms: OTUD4, DUBA6, HIN1, HIV-1 induced protein HIN-1, HIV-1-induced protein HIN-1, HSHIN1, KIAA1046, OTU domain containing 4
Gene ID:	54726
UniProt:	Q01804

Application Details

Application Notes:	Approved: ELISA, ICC, IHC, IHC-P (5 µg/mL), WB (0.25 - 0.5 µg/mL) Usage: OTUD4 antibody can be used for detection of OTUD4 by Western blot at 0.25-0.5 µg/mL.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, 0.02 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	avoid freeze thaw cycles. Store undiluted.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for 3 months and -20°C, stable for up to 1 year. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months



Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)

Image 1. Human Pancreas: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunohistochemistry

Image 2. Anti-OTUD4 antibody IHC of human pancreas. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.