



[Go to Product page](#)

Datasheet for ABIN609653

anti-CD8B antibody

2 Images

9 Publications

Overview

Quantity:	0.1 mg
Target:	CD8B
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD8B antibody is un-conjugated
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	CD8 positive Wistar rat splenic T cell hybridomas
Clone:	341
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 341 (also known as 34.1) recognizes rat CD8b, the 32-34 kDa beta chain of the CD8 coreceptor (extracellular epitope), expressed on T cell subsets and some other cell types, such as macrophages.
Cross-Reactivity (Details):	Rat
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	CD8B
Alternative Name:	CD8b (CD8B Products)
Background:	CD8b molecule,The CD8b (CD8 beta) subunit of CD8 T cell coreceptor is expressed in CD8 alpha/beta heterodimers on majority of MHC I-restricted conventional T cells and thymocytes and in CD8 alpha/alpha homodimers on subsets of memory T cells, intraepithelial lymphocytes, NK cells, macrophages, mast cells, and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 alpha/beta but not alpha/alpha dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.,Ly-3, Ly-C, Lyt-3
Gene ID:	926
UniProt:	P10966
Pathways:	TCR Signaling

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1.5 µg/mL.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Publications

Product cited in: Rossi, Del Principe, Rossi, Irno Consalvo, Luciano, Zucchetto, Bulian, Bomben, Dal Bo, Fangazio, Benedetti, Degan, Gaidano, Del Poeta, Gattei: "Prognostic impact of ZAP-70 expression in chronic lymphocytic leukemia: mean fluorescence intensity T/B ratio versus percentage of positive cells." in: **Journal of translational medicine**, Vol. 8, pp. 23, (2010) ([PubMed](#)).

Vroblova, Vrbacky, Hrudkova, Jankovicova, Schmitzova, Maly, Krejsek, Smolej: "Significant change in ZAP-70 expression during the course of chronic lymphocytic leukemia." in: **European journal of haematology**, Vol. 84, Issue 6, pp. 513-7, (2010) ([PubMed](#)).


Mabarrack, Turner, Mayrhofer: "Recent thymic origin, differentiation, and turnover of regulatory T cells." in: **Journal of leukocyte biology**, Vol. 84, Issue 5, pp. 1287-97, (2008) ([PubMed](#)).

Nohara, Kulka, Déry, Wills, Hirji, Gilchrist, Befus: "Regulation of CD8 expression in mast cells by exogenous or endogenous nitric oxide." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 167, Issue 10, pp. 5935-9, (2001) ([PubMed](#)).

Ellerman, Like: "Islet cell membrane antigens activate diabetogenic CD4+ T-cells in the BB/Wor rat." in: **Diabetes**, Vol. 48, Issue 5, pp. 975-82, (1999) ([PubMed](#)).


There are more publications referencing this product on: [Product page](#)

Images



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of rat splenocytes suspension stained using anti-rat CD8b (341) purified antibody (concentration in sample 1 µg/mL) GAM PE.



Flow Cytometry

Image 2. Surface staining of rat splenocytes using anti-CD8b (341) purified, GAM-APC.