

Datasheet for ABIN611634

anti-CD28 antibody**2** Images**7** Publications[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	CD28
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD28 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP)

Product Details

Immunogen:	Mouse A20J B lymphoma cells transfected with rat CD28
Clone:	JJ319
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody JJ319 reacts with an extracellular epitope of CD28, a disulfide-linked homodimeric type I glycoprotein (monomer of Mw 44 kDa) which is a critical costimulatory receptor of T cells.
Cross-Reactivity (Details):	Rat
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	CD28
Alternative Name:	CD28 (CD28 Products)
Background:	CD28 Molecule,CD28 is the critical T cell costimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting cells. Besides its costimulation role CD28 functions in preventing T cells from anergic hyporesponsive state or from undergoing premature apoptotic cell death. CD28 is also expressed on human fetal NK cells and some NK cell lines, whereas on murine NK cells the CD28 expression is much broader.,Tp44
Gene ID:	25660
UniProt:	P31042
Pathways:	TCR Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Production of Molecular Mediator of Immune Response

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 2-5 µg/mL.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Publications

Product cited in:	Kerstan, Armbruster, Leverkus, Hünig: "Cyclosporin A abolishes CD28-mediated resistance to
-------------------	--

CD95-induced apoptosis via superinduction of caspase-3." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 177, Issue 11, pp. 7689-97, (2006) ([PubMed](#)).

van den Brandt, Wang, Reichardt: "Resistance of single-positive thymocytes to glucocorticoid-induced apoptosis is mediated by CD28 signaling." in: **Molecular endocrinology (Baltimore, Md.)**, Vol. 18, Issue 3, pp. 687-95, (2004) ([PubMed](#)).


Lühder, Huang, Dennehy, Guntermann, Müller, Winkler, Kerkau, Ikemizu, Davis, Hanke, Hünig: "Topological requirements and signaling properties of T cell-activating, anti-CD28 antibody superagonists." in: **The Journal of experimental medicine**, Vol. 197, Issue 8, pp. 955-66, (2003) ([PubMed](#)).

Laskowski, Pratschke, Wilhelm, Dong, Beato, Taal, Gasser, Hancock, Sayegh, Tilney: "Anti-CD28 monoclonal antibody therapy prevents chronic rejection of renal allografts in rats." in: **Journal of the American Society of Nephrology : JASN**, Vol. 13, Issue 2, pp. 519-27, (2002) ([PubMed](#)).

Haspot, Villemain, Laflamme, Coulon, Olive, Tiollier, Souillou, Vanhove: "Differential effect of CD28 versus B7 blockade on direct pathway of allorecognition and self-restricted responses." in: **Blood**, Vol. 99, Issue 6, pp. 2228-34, (2002) ([PubMed](#)).


There are more publications referencing this product on: [Product page](#)

Images



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of rat splenocytes suspension stained using anti-rat CD28 (JJ319) purified antibody (concentration in sample 4 µg/mL) GAM APC.



Flow Cytometry

Image 2. Separation of rat CD28 positive splenocytes (red-filled) from CD28 negative splenocytes (black-dashed) in flow cytometry analysis (surface staining) of rat splenocytes suspension stained using anti-rat CD28 (JJ319) purified antibody (concentration in sample 4 µg/mL) GAM APC.