

Datasheet for ABIN612513

Rabbit anti-Goat IgG (Heavy & Light Chain) Antibody (DyLight 650)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Goat
Host:	Rabbit
Conjugate:	DyLight 650
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Immunogen:	Purified goat IgG, whole molecule
Characteristics:	Rabbit anti-goat IgG (H&L) - Affinity Pure, DyLight 650 Conjugate. Fluorophore: DyLight 650 (Ex = 652 nm, Em = 672 nm). Fluor Protein Ratio: Moles DyLight 650 per Mole Antibody.
Purification:	Affinity purified using solid phase mouse IgG (H&L)
Purity:	> 95 % based on SDS-PAGE

Target Details

Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody

Application Details

Application Notes:	<p>This conjugate is suitable for immunomicroscopy, flow cytometry.</p> <p>The optimal working dilution should be determined by the investigator. Suggested starting dilution: 1:20 - 1:2,000 for most applications</p>
Comment:	<p>Country of Origin: Rabbit serum was obtained from healthy animals of US origin, under the care of a registered veterinarian.</p> <p>DyLight is a trademark of Thermo Fisher Scientific, Inc. and its subsidiaries.</p>
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	1.0 mg/mL
Buffer:	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	<p>WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.</p>
Storage:	4 °C