



Datasheet for ABIN6142892  
**anti-KIR3DS1 antibody (AA 80-340)**



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	KIR3DS1
Binding Specificity:	AA 80-340
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIR3DS1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 80-340 of human KIR3DS1 (NP_001077008.1).
Sequence:	FNMSPVTTAH AGNYTCRGSHPHSPTGWSAP SNPMVIMVTG NHRKPSLLAH PGPLVKSGER VILQCWSDIM FEHFFLHKWEV ISKDPSRLVG QIHDGVSKAN FSI GMMRAL AGTYRCYGSV THTPYQLSAP SDPLDIVVTG LYEKPSLSAQ PGPKVQAGES VTLSCSSRSS YDMYHLSREG GAHERRLPAV RKNRRTFQAD FPLGPATHGG TYRCFGSFRH SPYEWSDPSD PLLVSVTGNP SSSWPSPTEP SSKSGNLRHL H
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Polyclonal Antibodies

## Target Details

---

Target:	KIR3DS1
Alternative Name:	KIR3DS1 ( <a href="#">KIR3DS1 Products</a> )
Background:	<p>Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several 'framework' genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules, thus, KIR proteins are thought to play an important role in regulation of the immune response. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.,KIR3DS1,CD158E2,KIR-123FM,KIR-G1,NKAT-10,NKAT10,KIR3DS1</p>
Molecular Weight:	42 kDa
Gene ID:	3813
UniProt:	<a href="#">Q14943</a>

## Application Details

---

Application Notes:	WB,1:500 - 1:2000,IHC,1:50 - 1:200
Comment:	HIGH QUALITY
Restrictions:	For Research Use only

## Handling

---


Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Handling

Storage: -20 °C


Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.

## Images




### Western Blotting

**Image 1.** Western blot analysis of extracts of various cell lines, using KIR3DS1 antibody.



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 2.** Immunohistochemistry of paraffin-embedded human liver using KIR3DS1 antibody.



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 3.** Immunohistochemistry of paraffin-embedded human mammary cancer using KIR3DS1 antibody.