antibodies - online.com












| Overview | |
|----------------------|---|
| Quantity: | 100 μL |
| Target: | MXI1 |
| Binding Specificity: | AA 170-295 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MXI1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 170-295 of human MXI1 (NP_569157.2). |
| Sequence: | HTTLGLLNKA KAHIKKLEEA ERKSQHQLEN LEREQRFLKW RLEQLQGPQE MERIRMDSIG STISSDRSDS EREEIEVDVE STEFSHGEVD NISTTSISDI DDHSSLPSIG SDEGYSSASV KLSFTS |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Characteristics: | Polyclonal Antibodies |
| Target Details | |
| Target: | MXI1 |

Target Details

| - | |
|---------------------|---|
| Alternative Name: | MXI1 (MXI1 Products) |
| Background: | Expression of the c-myc gene, which produces an oncogenic transcription factor, is tightly |
| | regulated in normal cells but is frequently deregulated in human cancers. The protein encoded |
| | by this gene is a transcriptional repressor thought to negatively regulate MYC function, and is |
| | therefore a potential tumor suppressor. This protein inhibits the transcriptional activity of MYC |
| | by competing for MAX, another basic helix-loop-helix protein that binds to MYC and is required |
| | for its function. Defects in this gene are frequently found in patients with prostate tumors. |
| | Three alternatively spliced transcripts encoding different isoforms have been described. |
| | Additional alternatively spliced transcripts may exist but the products of these transcripts have |
| | not been verified experimentally.,MXI1,MAD2,MXD2,MXI,bHLHc11,Epigenetics & Nuclear |
| | Signaling, Transcription Factors, MXI1 |
| Molecular Weight: | 20 kDa/21 kDa/26 kDa/32 kDa |
| Gene ID: | 4601 |
| UniProt: | P50539 |
| Pathways: | Maintenance of Protein Location |
| Application Details | |
| Application Notes: | WB,1:500 - 1:2000 |
| Comment: | HIGH QUALITY |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| | should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Avoid freeze / thaw cycles. |
| 3 | , |



Western Blotting

Image 1. Western blot analysis of extracts of various cell lines, using MXI1 antibody (ABIN6128218, ABIN6144206, ABIN6144208 and ABIN6215672) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 μg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.