



[Go to Product page](#)

Datasheet for ABIN6145194
anti-KIAA1274 antibody (AA 1-300)

2 Images

Overview

Quantity:	100 µL
Target:	KIAA1274 (PALD1)
Binding Specificity:	AA 1-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIAA1274 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human PALD1 (NP_055246.2).
Sequence:	MGTTASTAQQ TVSAGTPFEG LQSGTMDSR HSVSIHSFQS TSLHNSKAKS IIPNKVAPV ITYNCKEEFQ IHDELLKAHY TLGRLSDNTP EHYLVQGRYF LVRDVTEKMD VLGTVGSCGA PNFRQVQGGL TVFGMGQPSL SGFRRVLQKL QKDGHRECVI FCVREEPVLF LRADEDFVSY TPRDKQNLHE NLQGLGPGVR VESLELAIRK EIHDFAQLSE NTYHVYHNTD DLWGEPHAVA IHGEDDLHVT EEVYKRPLFL QPTYRYHRLP LPEQGSPLA QLDAFVSVLR ETPSLLQLRD
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Polyclonal Antibodies

Target Details


Target:	KIAA1274 (PALD1)
Alternative Name:	PALD1 (PALD1 Products)
Background:	PALD1,KIAA1274,PALD,paladin,Signal Transduction,Cell Biology & Developmental Biology,Growth factor,PALD1
Molecular Weight:	96 kDa
Gene ID:	27143
UniProt:	Q9ULE6

Application Details

Application Notes:	WB,1:500 - 1:2000,IF,1:50 - 1:100
Comment:	HIGH QUALITY
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.



Immunofluorescence

Image 1. Immunofluorescence analysis of L929 cells using P Polyclonal Antibody (ABIN6133270, ABIN6145194, ABIN6145195 and ABIN6216692) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using P antibody (ABIN6133270, ABIN6145194, ABIN6145195 and ABIN6216692) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.