



[Go to Product page](#)

Datasheet for ABIN6149504  
**anti-TRIM15 antibody (AA 1-330)**

1 Image

Overview

Quantity:	100 µL
Target:	TRIM15
Binding Specificity:	AA 1-330
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM15 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-330 of human TRIM15 (NP_150232.2).
Sequence:	MPATPSLKVV HELPACTLCA GPLEDAVTIP CGHTFCRLCL PALSQMGAQS SGKILLCPLC QEEEAETPM APVPLGPLGE TYCEEHGEKI YFFCENDAEF LCVFCREGPT HQAHTVGFLD EAIQPYRDRL RSRLEALSTE RDEIEDVKCQ EDQKLQVLLT QIESKKHQVE TAFERLQQEL EQQRCLLLAR LRELEQQIWK ERDEYITKVS EEVTRLGAQV KELEEKCCQP ASELLQDVRV NQSRCEMKTf VSPEAISPDL VKKIRDFHRK ILTLPEMMRM FSENLAHHLE IDSGVITLDP QTASRSLVLS EDRKSVRYTR QKKSPLDSPL
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Characteristics:	Polyclonal Antibodies

## Target Details

---

Target:	TRIM15
Alternative Name:	TRIM15 ( <a href="#">TRIM15 Products</a> )
Background:	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to the cytoplasm. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.,TRIM15,RNF93,ZNF178,ZNF7,Epigenetics & Nuclear Signaling,Cell Biology & Developmental Biology,Ubiquitin,TRIM15
Molecular Weight:	12 kDa/52 kDa
Gene ID:	89870
UniProt:	<a href="#">Q9C019</a>

## Application Details


---

Application Notes:	WB,1:500 - 1:2000
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.



### Western Blotting

**Image 1.** Western blot analysis of extracts of various cell lines, using TRIM15 antibody (ABIN6132733, ABIN6149504, ABIN6149505 and ABIN6215428) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.