

Datasheet for ABIN6150086
anti-VEGFA antibody (AA 1-147)

6 Images

4 Publications

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	VEGFA
Binding Specificity:	AA 1-147
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VEGFA antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-147 of human VEGF (NP_001165099.1).
Sequence:	MNLLSWVHW SLALLYLHH AKWSQAAPMA EGGGQNHHEV VKFMDVYQRS YCHPIETLVD IFQEYPDEIE YIFKPSCVPL MRCGGCCNDE GLECVPTES NITMQIMRIK PHQGQHIGEM SFLQHNKCEC RPKKDRARQE KCDKPRR
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Polyclonal Antibodies

Target Details

Target:	VEGFA
---------	-------

Target Details

Alternative Name:	VEGFA (VEGFA Products)
Background:	<p>This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site.,VEGFA,MVCD1,VEGF,VPF,L VEGFA,VEGF A,Cancer,Invasion and Metastasis,Signal Transduction,Cell Biology & Developmental Biology,Apoptosis,Growth factor,VEGF,Endocrine & Metabolism,Immunology & Inflammation,Cytokines,Cardiovascular,Angiogenesis,Angiogenic growth factors,Hypoxia,VEGFA</p>
Molecular Weight:	15-27 kDa/34-45 kDa
Gene ID:	7422
UniProt:	P15692
Pathways:	RTK Signaling , Glycosaminoglycan Metabolic Process , Regulation of Cell Size , Tube Formation , Signaling Events mediated by VEGFR1 and VEGFR2 , Platelet-derived growth Factor Receptor Signaling , VEGFR1 Specific Signals , VEGF Signaling

Application Details


Application Notes:	WB,1:500 - 1:2000
Comment:	HIGH QUALITY
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.


Publications

Product cited in:	<p>Huang, Tong, Rehman, Wang, Zhang, Wang, Li, Yang: "Oxygen Supplementation Ameliorates Tibial Development via Stimulating Vascularization in Tibetan Chickens at High Altitudes." in: International journal of biological sciences, Vol. 13, Issue 12, pp. 1547-1559, (2018) (PubMed).</p> <p>Lin, Cao, Wang, Hu, Liu, Li, Chen, Li, Liu, Wang, Zheng: "Long non-coding RNA UBE2CP3 enhances HCC cell secretion of VEGFA and promotes angiogenesis by activating ERK1/2/HIF-1α/VEGFA signalling in hepatocellular carcinoma." in: Journal of experimental & clinical cancer research : CR, Vol. 37, Issue 1, pp. 113, (2018) (PubMed).</p> <p>Huang, Rehman, Lan, Qiu, Zhang, Iqbal, Luo, Mehmood, Zhang, Li: "Tibial dyschondroplasia is highly associated with suppression of tibial angiogenesis through regulating the HIF-1α/VEGF/VEGFR signaling pathway in chickens." in: Scientific reports, Vol. 7, Issue 1, pp. 9089, (2017) (PubMed).</p> <p>Yin, Zhao, Li, Yan, Zhou, Chen, Wang: "miR-320a mediates doxorubicin-induced cardiotoxicity by targeting VEGF signal pathway." in: Aging, Vol. 8, Issue 1, pp. 192-207, (2016) (PubMed).</p>
-------------------	--




Immunofluorescence

Image 1. Immunofluorescence analysis of HeLa cells using VEGFA Rabbit pAb (ABIN6128908, ABIN6150086, ABIN6150088 and ABIN6221373) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Western Blotting

Image 2. Western Blot Analysis using ABIN6150086



Western Blotting

Image 3. Western blot analysis of extracts of H460 cells, using antibody at 1:1000 dilution.

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN6150086.