

Datasheet for ABIN6157201 **anti-Melanoma Marker antibody (CF®594)**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Melanoma Marker
Reactivity:	Human, Dog, Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Melanoma Marker antibody is conjugated to CF®594
Application:	Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Purpose:	Mouse Monoclonal anti-Melanoma Marker (PNL2), CF594 Conjugate
Immunogen:	Melanocyte antigen
Clone:	PNL2
Isotype:	IgG1

Characteristics: Anti-PNL2 is a novel monoclonal antibody, which has recently been introduced as an immunohistochemical reagent to stain melanocytes and tumors derived therefrom. The antigen recognized by PNL2 is different from Melan A and gp100. Its epitope is not destroyed by digestion with neuraminidase i.e. its epitope is not glycosylated. Anti-PNL2 may be most useful because of its high sensitivity for metastatic melanoma (87 %), as opposed to 76 % for anti-HMB45 and 82 % for anti-MART-1. Anti-PNL2 labels intra-epidermal nevi while the dermal component of compound nevi are largely non-reactive with anti-PNL2. Antibodies against PNL2, MART-1 (Melan A) and HMB45 stain most clear cell sarcoma cells and a few cells in angio-

Product Details

myolipomas and lymphangioleiomyomatosis. Anti-PNL2 is a useful antibody for the identification of melanomas and clear cell sarcomas. Differential diagnosis is aided by the results from a panel of antibodies, including antibodies against HMB45, MART-1, tyrosinase, and MiTF. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the < a href="https://biotium.com/wp-content/uploads/2013/07/CF-dye-selection-guide.pdf" target="_blank" rel="noopener">CF® Dye Selection Guide for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Target:	Melanoma Marker
Abstract:	Melanoma Marker Products
Background:	Human Melanoma Associated Antigen, PNL2
Molecular Weight:	Multiple (160 kDa, 100 kDa and <10 kDa)

Application Details

Application Notes:	Immunohistology (formalin) <ul style="list-style-type: none">• Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes• Immunofluorescence 0.5-1 µg/mL• Optimal dilution for a specific application should be determined by user
Comment:	Melanoma
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	100 µg/mL
Buffer:	PBS/0.1 % BSA/0.05 % azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

should be handled by trained staff only.

Handling Advice: Protect from light