

Datasheet for ABIN6168087

anti-Uroplakin 3A antibody (AA 260-279) (CF®640R)



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	Uroplakin 3A (UPK3A)	
Binding Specificity:	AA 260-279	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Uroplakin 3A antibody is conjugated to CF®640R	
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS)	

Product Details

Purpose:	Polyclonal-anti-UPK3A (Uroplakin-3A) (Rabbit-PAb), CF640R Conjugate	
Immunogen:	A synthetic peptide, aa260-279 (GASESSYTSVNRGPPLDRAE) of human UPK3A	
Clone:	PAb	
Isotype:	IgG	
Characteristics:	Uroplakin-3A (UPK3A) is a component of the asymmetric unit membrane (AUM). It is a highly	

Uroplakin-3A (UPK3A) is a component of the asymmetric unit membrane (AUM). It is a highly specialized bio-membrane made by terminally differentiated urothelial cells. The protein may play an important role in AUM-cytoskeleton interaction in terminally differentiated urothelial cells. UPK3A also contributes to the formation of urothelial glycocalyx, which may play an important role in preventing bacterial adherence through FimH bacterial protein binding leading to bladder infection. UPK3A has been shown as a helpful marker for the detection of bladder cancer. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes

and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Target:	Uroplakin 3A (UPK3A)	
Alternative Name:	-UPK3A (Uroplakin-3A) (-) (UPK3A Products)	
Background:	UP3a, UPIIIA, UPK3A, Uroplakin III (UPIII), Uroplakin3A	
Molecular Weight:	~30 kDa	
Gene ID:	7380, 632787	
UniProt:	075631	

Application Details

Application Notes:	Flow Cytometry 0.5-1 µg/million cells/0.1 mL
	 Immunofluorescence 1-2 μg/mL Western blotting 0.5-1 μg/mL Optimal dilution for a specific application should be determined by user
Comment:	HEK293 cells or bladder carcinoma
Restrictions:	For Research Use only

Handling

Format:	Liquid	
Concentration:	100 μg/mL	
Buffer:	PBS/0.1 % BSA/0.05 % azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Protect from light	