

Datasheet for ABIN6170506  
**anti-CD45RB antibody (CF®647)**



[Go to Product page](#)

## Overview

|              |                                                                            |
|--------------|----------------------------------------------------------------------------|
| Quantity:    | 100 µL                                                                     |
| Target:      | CD45RB                                                                     |
| Reactivity:  | Human                                                                      |
| Host:        | Mouse                                                                      |
| Clonality:   | Monoclonal                                                                 |
| Conjugate:   | This CD45RB antibody is conjugated to CF®647                               |
| Application: | Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (IHC) |

## Product Details

|            |                                                                             |
|------------|-----------------------------------------------------------------------------|
| Purpose:   | Mouse Monoclonal anti-CD45RB (BRA-11G), CF647 Conjugate                     |
| Immunogen: | Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V) |
| Clone:     | BRA-11G                                                                     |
| Isotype:   | IgG1 kappa                                                                  |

**Characteristics:** CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and

## Product Details

leukemias of B-cell origin. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

## Target Details

|                   |                                                                                                                                                                                             |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target:           | CD45RB                                                                                                                                                                                      |
| Alternative Name: | CD45RB ( <a href="#">CD45RB Products</a> )                                                                                                                                                  |
| Background:       | B220, CD45R, GP180, Leukocyte common antigen (LCA), Loc, Ly-5, Lyt-4, Protein tyrosine phosphatase receptor type C (PTPRC), Receptor-type tyrosine-protein phosphatase C, T200 glycoprotein |
| Molecular Weight: | 180-220 kDa                                                                                                                                                                                 |
| Gene ID:          | 5788, 654514                                                                                                                                                                                |
| UniProt:          | <a href="#">P08575</a>                                                                                                                                                                      |

## Application Details

|                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | Does not react with rat or rhesus monkey, others not known <ul style="list-style-type: none"><li>• Immunohistology (formalin)</li><li>• Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes</li><li>• Immunofluorescence 0.5-1 µg/mL</li><li>• Flow Cytometry 0.5-1 µg/million cells/0.1 mL</li><li>• Optimal dilution for a specific application should be determined by user</li></ul> |
| Comment:           | Ramos, U-698, or GA-10 cells. Tonsil                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Restrictions:      | For Research Use only                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

## Handling

|                |                            |
|----------------|----------------------------|
| Format:        | Liquid                     |
| Concentration: | 100 µg/mL                  |
| Buffer:        | PBS/0.1 % BSA/0.05 % azide |
| Preservative:  | Sodium azide               |

## Handling

---

|                    |                                                                                                                        |
|--------------------|------------------------------------------------------------------------------------------------------------------------|
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
|--------------------|------------------------------------------------------------------------------------------------------------------------|

---

|                  |                    |
|------------------|--------------------|
| Handling Advice: | Protect from light |
|------------------|--------------------|