

Datasheet for ABIN6170612
anti-CD45RB antibody (CF®405S)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CD45RB
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD45RB antibody is conjugated to CF®405S
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (IHC), Western Blotting (WB)

Product Details

Purpose:	Mouse Monoclonal anti-CD45RB (PTPRC/1132), CF405S Conjugate
Immunogen:	Recombinant full-length human CD45RB protein
Clone:	PTPRC-1132
Isotype:	IgG1 kappa
Characteristics:	CD45R, also designated CD45 and PTPRC, is identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature

Product Details

B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Target:	CD45RB
Alternative Name:	CD45RB (CD45RB Products)
Background:	B220, CD45R, GP180, Leukocyte common antigen (LCA), Loc, Ly-5, Lyt-4, Protein tyrosine phosphatase receptor type C (PTPRC), Receptor-type tyrosine-protein phosphatase C, T200 glycoprotein
Molecular Weight:	180-220 kDa
Gene ID:	5788, 654514
UniProt:	P08575

Application Details

Application Notes:	Does not react with rat or monkey, others not known • Immunohistology (formalin) • Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes • Immunofluorescence 0.5-1 µg/mL • Flow Cytometry 0.5-1 µg/million cells/0.1 mL • Western blotting 0.5-1.0 µg/mL • Optimal dilution for a specific application should be determined by user
Comment:	Ramos, U-698, or GA-10 cells. Tonsil
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	100 µg/mL

Handling

Buffer:	PBS/0.1 % BSA/0.05 % azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from light