

Datasheet for ABIN6174606

anti-CFTR antibody (CF®640R)



Overview

Quantity:	100 μL
Target:	CFTR
Reactivity:	Human, Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CFTR antibody is conjugated to CF®640R
Application:	Immunofluorescence (IF), Immunohistochemistry (IHC), Immunostaining (ISt), Western Blotting (WB)

Product Details

Characteristics	This antibody recognizes a protein of 16E 170 kDs identified as evetic fibracia transmembrane		
Isotype:	lgG2a		
Clone:	CFTR-1342		
Immunogen:	Recombinant human full-length CFTR		
Purpose:	Mouse Monoclonal anti-CFTR CFTR/1342), CF640R Conjugate		

Characteristics:

This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3',5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of

epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Target:	CFTR
Alternative Name:	CFTR (CFTR Products)
Molecular Weight:	165-170 kDa
Gene ID:	1080

Application Details

lication	

Immunohistology (formalin) 0.5-1 μg/mL

- Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM
 EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min
- Immunofluorescence 0.5-1 µg/mL
- Western blotting 0.5-1 μg/mL
- · Optimal dilution for a specific application should be determined by user

Comment:

MOLT-4 cells. Pancreas, Kidney or Placenta.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	100 μg/mL
Buffer:	PBS/0.1 % BSA/0.05 % azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

		1 .
_	コヘロム	lino
	land	11111

Handling Advice:

Protect from light