

[Go to Product page](#)

Datasheet for ABIN6178766

**Recombinant anti-Keratin Acidic (AE1) antibody (CF®488A)**

## Overview

Quantity:	100 µL
Target:	Keratin Acidic (AE1)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	CF®488A
Application:	Immunohistochemistry (IHC), Immunofluorescence (IF), Western Blotting (WB), Flow Cytometry (FACS)

## Product Details

Purpose:	Recombinant Rabbit Monoclonal anti-Cytokeratin, Acidic (KRTL/1577R), CF488A Conjugate
Immunogen:	Recombinant human KRT77 protein fragment
Clone:	KRTL-1577R
Isotype:	IgG kappa
Characteristics:	<p>This MAb recognizes the 56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI &lt;5.7) and basic (pI &gt;6.0) subfamilies. The acidic keratins have molecular weights (MW) of 56.5, 55, 51, 50, 50, 48, 46, 45, and 40 kDa. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. Primary antibodies are available purified, or with a selection of fluorescent</p>

## Product Details

CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

## Target Details

Target:	Keratin Acidic (AE1)
Alternative Name:	Cytokeratin, Acidic ( <a href="#">AE1 Products</a> )
Background:	K1B, KRT1B, K77, CK-1B, Keratin 1B, Keratin-77, Cytokeratin-1B
Molecular Weight:	56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19)
Gene ID:	374454, 334989
UniProt:	<a href="#">Q7Z794</a>

## Application Details

Application Notes:	Immunohistology (formalin) 1-2 µg/mL <ul style="list-style-type: none"><li>Immunofluorescence 1-2 µg/mL Western blotting 0.5-1 µg/mL</li><li>Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min</li><li>Flow Cytometry 0.5-1 µg/million cells/0.1 mL</li><li>Optimal dilution for a specific application should be determined by user</li></ul>
Comment:	Skin, Squamous cell carcinoma (SCC)
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	100 µg/mL
Buffer:	PBS/0.1 % BSA/0.05 % azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from light