

Datasheet for ABIN6180871 **anti-HCG beta antibody**



[Go to Product page](#)

Overview

Quantity:	50 µL
Target:	HCG beta
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HCG beta antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Purpose:	Mouse Monoclonal anti-HCG beta (HCGb/54+HCGb/459), Purified, BSA-free
Immunogen:	Recombinant hCG beta protein (HCGb/54 and HCGb/459)
Clone:	HCGb-54 HCGb-459
Isotype:	IgG
Characteristics:	This MAb reacts with a protein of 22 kDa, identified as β sub-unit of HCG. It does not cross react with the α sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the α and β subunits. The α subunit is identical to that of thyroid stimulating hormone (TSH), follicle stimulating hormone (FSH), and luteinizing hormone (LH). hCG MAb detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate anti-hCG positivity in 90 % and

Product Details

60 % of cases respectively. 20 % of lung squamous cell carcinomas are positive. hCG expression by non-trophoblastic tumors may indicate aggressive behavior. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Purification: Purified

Target Details

Target: HCG beta

Abstract: [HCG beta Products](#)

Background: CG-beta, CGB3, CGB5, CGB7, CGB8, Choriogonadotropin Subunit beta, hCGB

Molecular Weight: 22 kDa

Gene ID: 1082, 172944

UniProt: [P01233](#)

Application Details

Application Notes: Immunohistology formalin-fixed 0.5-1 µg/mL

- Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes
- Optimal dilution for a specific application should be determined by user

Comment: Placenta

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS (no BSA, no azide)

Preservative: Azide free