antibodies .- online.com





anti-LGALS13 antibody (AA 23-134) (CF®640R)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	LGALS13
Binding Specificity:	AA 23-134
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This LGALS13 antibody is conjugated to CF®640R
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (IHC), ELISA, Coating (Coat)

Product Details

Purpose:	Mouse Monoclonal anti-Galectin-13 / Placental-Protein-13 (PP13/1162), CF640R Conjugate
Immunogen:	Recombinant human Galectin-13 protein fragment (aa23-134) (exact sequence is proprietary)
Clone:	PP13-1162
Isotype:	lgG2b
Characteristics:	This antibody recognizes a 32 kDa protein, which is identified as homodimer of galectin-13
	(also known as PP13). Galectins are a family of soluble ?-galactoside-binding lectins that
	modulate cell-to-cell adhesion and cell-to-extracellular matrix (ECM) interactions and play a role
	in tumor progression, pre-mRNA splicing and apoptosis. One such member, galectin-13, is a

139 amino acid protein that contains one galectin domain. Possessing lysophospholipase

activity, galectin-13 exists as a disulfide-linked homodimer. Galectin-13 is suggested to have a

developmental role in the placenta and may display immunobiological activity in feto-maternal blood-spaces. Screening for galectin-13 in maternal serum during the first trimester of pregnancy may serve as a diagnostic marker to predict preterm pre-eclampsia. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Target Details

Target:	LGALS13
Alternative Name:	Galectin-13 / Placental-Protein-13 (LGALS13 Products)
Background:	Gal-13, GAL13, Galactoside-binding soluble lectin 13, Galectin-13, Galectin13, Lectin galactoside binding soluble 13, LGALS13, PLAC8, Placental protein 13 (PP13), Placental tissue protein 13
Molecular Weight:	16 kDa (monomer), 32 kDa (homodimer)
Gene ID:	29124, 23671
UniProt:	Q9UHV8

Application Details

Ap	рп	cat	ion	INO	tes:

Immunohistology (formalin) 0.5-1 μg/mL

- Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM
 EDTA pH 9.0 or 10 mM citrate buffer pH 6.0 for 10-20 min followed by cooling at RT for 20 min
- Immunofluorescence 0.5-1 µg/mL
- Flow Cytometry 0.5-1 μg/million cells/0.1 mL
- · ELISA For coating, order Ab without BSA
- · Optimal dilution for a specific application should be determined by user

Comment:

K-562 cells. Placenta or Spleen.

Restrictions:

For Research Use only

Handling

Format: Liquid

Handling

Concentration:	100 μg/mL
Buffer:	PBS/0.1 % BSA/0.05 % azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect from light