

Datasheet for ABIN6189953

anti-TCP1 alpha/CCTA antibody (AA 3-106)[Go to Product page](#)

Overview

Quantity:	50 µL
Target:	TCP1 alpha/CCTA (TCP1)
Binding Specificity:	AA 3-106
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TCP1 alpha/CCTA antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS)

Product Details

Purpose:	Mouse Monoclonal anti-p63 (TP63/1786), Purified, BSA-free
Immunogen:	Recombinant human p63 protein fragment (aa3-106) (exact sequence is proprietary)
Clone:	TP63-1786
Isotype:	IgG2b kappa
Characteristics:	p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal prostate glands, however, it was not expressed in malignant tumors of the prostate. As a result, p63 has been reported as a useful marker for differentiating benign from malignant lesions in the prostate, particularly when used in combination with markers of high molecular weight cytokeratins and the prostate-specific marker AMACR (P504S). p63 has also been shown to be a sensitive marker for lung squamous

Product Details

cell carcinomas (SqCC), with a sensitivity of ~90 % . Specificity for lung SqCC, vs. lung adenocarcinoma (LADC), is approximately 80 %. In breast tissue, p63 has been identified in myoepithelial cells of normal ducts. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Purification: Purified

Target Details

Target: TCP1 alpha/CCTA (TCP1)

Alternative Name: p63 (TCP1 Products)

Molecular Weight: 63 kDa

Gene ID: 8626, 137569

UniProt: [Q9H3D4](#)

Application Details

Application Notes: Flow Cytometry 0.5-1 µg/million cells/0.1 mL

- Immunofluorescence 1-2 µg/mL
- Western blotting 0.5-1 µg/mL
- Optimal dilution for a specific application should be determined by user

Comment: HEK293 cells or Prostate Carcinoma or Lung or bladder squamous cell carcinoma

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS (no BSA, no azide)

Preservative: Azide free