

Datasheet for ABIN619900 anti-DHAR2 antibody



Overview

Quantity:	200 µg
Target:	DHAR2
Reactivity:	Arabidopsis thaliana
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide derived from known DHAR1 sequence of Arabidopsis
	thaliana Q9FRL8, At1g75270
Cross-Reactivity (Details):	Not reactive in: no confirmed exceptions from predicted reactivity are currently known
Predicted Reactivity:	dicots including: Ricinus communis, trees: Populus trichocarpa
Characteristics:	Expected / apparent Molecular Weight of the Antigene: 23.6 / 23.4 kDa
Purification:	serum

Target Details

Target:	DHAR2
Alternative Name:	Dehydroascorbate Reductase 2 (DHAR2) (DHAR2 Products)
Background:	AGI Code: At1g75270 DHAR2 (Dehydroascorbate Reductase 2) the protein is induced by jasmonic acid and oxidative
	chemical stresses and is a key component of the ascorbate recycling system. Involved in redox

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN619900 | 05/14/2025 | Copyright antibodies-online. All rights reserved.

Target Details	
	homeostasis under biotic and abiotic inducers. Localized in cytoplasm. Synonymes: chloride intracellular channel homolog 2, CLIC homolog 2, glutathione-dependent dehydroascorbate reductase 2, DHAR2, CytDHAR, GSH-dependent dehydroascorbate reductase 2.
Molecular Weight:	expected: 23.6 kDa, apparent: 23.4 kDa
UniProt:	Q9FRL8
Application Details	
Application Notes:	1 : 5000 with standard ECL (WB)
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	For reconstitution add 200 μL of sterile water.
Storage:	-20 °C
Storage Comment:	store lyophilized/reconstituted at -20°C, once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.