

Datasheet for ABIN622080 TMPRSS15 Protein



Overview Quantity: 1000 IU Target: TMPRSS15 Origin: Pig Source: Pig Protein Type: Native **Product Details** Characteristics: Native Porcine Enteropeptidase/Enterokinase **Target Details** TMPRSS15 Target: Alternative Name: Enteropeptidase/Enterokinase (TMPRSS15 Products) Porcine enteropeptidase is a specific protease which cleaves after the lysine at its recognition Background: site: Asp-Asp-Asp-Asp-Lys. Enterokinase will not cleave a site followed by proline. Theoretical Mw is 21,880 Dalton, the apparent Mw on SDS-PAGE is about 40 kDa. If a fusion tag is located in the N-terminus with an enterokinase site, enterokinase will be able to remove the fusion tag and to generate the protein exactly as you need without adding any unwanted residues. RayBiotechs enterokinase is a highly purified enterokinase from porcine. The enzyme has been extensively purified and tested to ensure that there are no other contaminating proteases. Introduction: Enteropeptidase or enterokinase is an enzyme involved in human digestion. It is produced by cells in the duodenum wall, and is secreted from duodenum's glands, the crypts of Lieberkühn, whenever ingested food enters the duodenum from the stomach.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN622080 | 07/26/2024 | Copyright antibodies-online. All rights reserved. Enteropeptidase has the critical job of turning trypsinogen (a zymogen) to trypsin, indirectly activating a number of pancreatic digestive enzymes. Enteropeptidase is a serine protease enzyme(EC3.4.21.9). Enteropeptidase is a part of the Chymotrypsin-clan of serine proteases, and is structurally similar to these proteins. Synonyms: Enteropeptidase, EC 3.4.21.9, Enterokinase, Serine protease 7, ENTK, MGC133046.

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	2 IU/µl, 50mM Tris-HCl, pH 8.0, 0.5M NaCl and 50% glycerol.
Storage:	-20 °C