

Datasheet for ABIN6239769
IL-20 Protein (AA 25-176) (His tag)

3 Images

[Go to Product page](#)

Overview

Quantity:	50 µg
Target:	IL-20 (IL20)
Protein Characteristics:	AA 25-176
Origin:	Human
Source:	Escherichia coli (E. coli)
Biological Activity:	Active
Purification tag / Conjugate:	This IL-20 protein is labelled with His tag.
Application:	Activity Assay (AcA), Cell Culture (CC)

Product Details

Characteristics:	Tag location: N-terminal His Tag
Purity:	> 90 %
Biological Activity Comment:	IL20 (Interleukin-20) is a cytokine structurally related to interleukin 10, which is produced by activated keratinocytes and monocytes. It is accepted that IL20 regulates proliferation and differentiation of keratinocytes during inflammation, particularly inflammation associated with the skin. Thus, proliferation assay of IL20 was conducted using ECV-304 cells. Briefly, ECV-304 cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard 1640 prior to the addition of various concentrations of IL20. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10µL of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-

Product Details

4 hours at 37°C. Proliferation of ECV-304 cells after incubation with IL20 for 48h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with human recombinant IL20 for 48h. The result was shown in Figure 2. It was obvious that human IL20 significantly decreased cell viability of ECV-304 cells.

Target Details


Target:	IL-20 (IL20)
Abstract:	IL20 Products
Background:	Alternative Names: IL10D, IL10-D, ZCYTO10, Cytokine Zcyto10
Molecular Weight:	21kDa
UniProt:	Q9NYY1
Pathways:	Protein targeting to Nucleus

Application Details

Application Notes:	Isoelectric Point: 8.9
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 1 mM DTT, 0.01 % SKL, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), Other preservative, ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.



SDS-PAGE

Image 1. Figure. SDS-PAGE; Sample: Active recombinant IL20, Human.




Figure 2. Cell proliferation of ECV-304 cells after stimulated with IL20.

Image 2. IL20 (Interleukin-20) is a cytokine structurally related to interleukin 10, which is produced by activated keratinocytes and monocytes. It is accepted that IL20 regulates proliferation and differentiation of keratinocytes during inflammation, particularly inflammation associated with the skin. Thus, proliferation assay of IL20 was conducted using ECV-304 cells. Briefly, ECV-304 cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard 1640 prior to the addition of various concentrations of IL20. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10µL of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. Proliferation of ECV-304 cells after incubation with IIL20 for 48h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with human recombinant IL20 for 48h. The result was shown in Figure 2. It was obvious that human IL20 significantly decreased cell viability of ECV-304 cells.




Figure 1. Cell proliferation of ECV-304 cells after stimulated with IL20.

(A) ECV-304 cells cultured in 1640, stimulated with 500ng/mL IL20 for 48h;

(B) Unstimulated ECV-304 cells cultured in 1640 for 48h.

Image 3.