

Datasheet for ABIN6239864

FGF22 Protein (AA 26-162) (His tag)**3** Images[Go to Product page](#)

Overview

Quantity:	50 µg
Target:	FGF22
Protein Characteristics:	AA 26-162
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Biological Activity:	Active
Purification tag / Conjugate:	This FGF22 protein is labelled with His tag.
Application:	Activity Assay (AcA), Cell Culture (CC)

Product Details

Characteristics:	Tag location: N-terminal His Tag
Purity:	> 95 %
Biological Activity Comment:	<p>FGF22 (Fibroblast growth factor 22) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. A proliferation assay was conducted to detect the bioactivity of recombinant mouse FGF22 using 3T3 cells. Briefly, 3T3 cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of FGF22. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10µL of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured</p>

Product Details

using a microplate reader after incubating the plate for 1-4 hours at 37°C. Proliferation of 3T3 cells after incubation with FGF22 for 48h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant FGF22 for 48h. The result was shown in Figure 2. It was obvious that FGF22 significantly increased cell viability of 3T3 cells.

Target Details

Target:	FGF22
Abstract:	FGF22 Products
Molecular Weight:	20kDa
UniProt:	Q9ESS2
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway

Application Details

Application Notes:	Isoelectric Point: 11.7
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 1 mM DTT, 0.01 % SKL, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), Other preservative, ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.




Image 1.




Image 2.




Figure 2. Cell proliferation of 3T3 cells after stimulated with FGF22.

Image 3. FGF22 (Fibroblast growth factor 22) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. A proliferation assay was conducted to detect the bioactivity of recombinant mouse FGF22 using 3T3 cells. Briefly, 3T3 cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of FGF22. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by

Cell Counting Kit-8 (CCK-8). Briefly, 10 μ L of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. *Proliferation of 3T3 cells after incubation with FGF22 for 48h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant FGF22 for 48h. The result was shown in Figure 2. It was obvious that FGF22 significantly increased cell viability of 3T3 cells.*