antibodies .- online.com







ARG Protein (AA 1-323) (His tag)



Image



Overview

Quantity:	50 μg
Target:	ARG
Protein Characteristics:	AA 1-323
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Biological Activity:	Active
Purification tag / Conjugate:	This ARG protein is labelled with His tag.
Application:	Activity Assay (AcA), Cell Culture (CC)

Product Details

Characteristics:	Tag location: N-terminal His Tag
Purity:	> 95 %
Biological Activity Comment:	Arginase (Arg) is an enzyme that catalyzes the degradation of arginine to produce urea and
	omithine, which is crucial in the urea cycle. In most mammals, two isozymes of this enzyme
	exist; the first, Arginase I, functions in the urea cycle, and is located primarily in the cytoplasm
	of the liver. The second isozyme, Arginase II, has been implicated in the regulation of the
	arginine/ornithine concentrations in the cell. Besides, Ubiquitin Carboxyl Terminal Hydrolase L5
	(UCHL5) has been identified as an interactor of Arg, thus a binding ELISA assay was conducted
	to detect the interaction of recombinant mouse Arg and recombinant mouse UCHL5. Briefly,
	Arg were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then
	transferred to UCHL5-coated microtiter wells and incubated for 2h at 37°C. Wells were washed

with PBST and incubated for 1h with anti-Arg pAb, then aspirated and washed 3 times. After

Product Details

incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50μ L stop solution to the wells and read at 450nm immediately. The binding activity of of Arg and UCHL5 was shown in Figure 1, and this effect was in a dose dependent manner The binding activity of Arg with UCHL5.

Target Details

Target:	ARG
Abstract:	ARG Products
Background:	Alternative Names: ARG1, Arginase I, Liver Arginase
Molecular Weight:	37kDa
UniProt:	Q61176

Application Details

Application Notes:	Isoelectric Point: 6.6
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 1 mM DTT, 0.01 % SKL, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), Other preservative, ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.




Image 1.