antibodies - online.com







anti-Macrophage Mannose Receptor 1 antibody (FITC)



Images



()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	1 V I	ew


Quantity:	100 tests	
Target:	Macrophage Mannose Receptor 1 (MRC1)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This Macrophage Mannose Receptor 1 antibody is conjugated to FITC	
Application:	Flow Cytometry (FACS)	

Product Details

Immunogen:	Purified human mannose receptor	
Clone:	15-2	
Isotype:	IgG1 kappa	
Specificity:	The mouse monoclonal antibody 15-2 (also known as MR15-2) recognizes an extracellular epitope of CD206 (macrophage mannose receptor, MMR), a 162-175 kDa type I transmembrane protein expressed mainly on macrophages, dendritic cells and hepatic or lymphatic endothelial cells, but not on monocytes.	
Cross-Reactivity (Details):	Human	
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.	


Target Details

Target:	Macrophage Mannose Receptor 1 (MRC1)	
Alternative Name:	CD206 (MRC1 Products)	
Background:	Mannose receptor C-type 1,CD206 (macrophage mannose receptor, MMR), also known as mannose receptor C1 (MRC1), is a type I transmembrane glycoprotein serving as pattern recognition receptor for carbogydrate groups on the surface of bacteria, fungi and other pathogens. Expressed mainly on tissue macrophages and dendritic cells, CD206 mediates endocytosis of these pathogens and presentation of their antigens to the adaptive immune system. CD206 can also be detected in a soluble form in human plasma and is elevated in patients with acute sepsis.,MMR, MRC1, CLEC13DL	
Gene ID:	4360	
UniProt:	P22897	
Application Details		
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.	
Comment:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.	
Restrictions:	For Research Use only	
Handling		
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.	
Storage:	4 °C	
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells stained using anti-human CD206 (15-2) FITC antibody (4 μ L reagent per milion cells in 100 μ L of cell suspension).



Flow Cytometry

Image 2. Flow cytometry multicolor surface staining of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells stained using anti-human CD206 (15-2) FITC antibody (4 μ L reagent per milion cells in 100 μ L of cell suspension) and anti-human CD11c (BU15) APC antibody (10 μ L reagent per milion cells in 100 μ L of cell suspension).



Flow Cytometry

Image 3. Separation of human CD206 positive CD11c positive dendritic cells differentiated upon monocyte stimulation (GM-CSF + IL-4) (red-filled) from non-stimulated lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human stimulated (GM-CSF + IL-4) peripheral blood mononuclear cells stained using antihuman CD206 (15-2) FITC antibody (4 μ L reagent / 100 μ L of peripheral whole blood).