

Datasheet for ABIN6242156

anti-Adenylate Kinase 2 antibody (AA 147-183)[Go to Product page](#)**4** Images

Overview

Quantity:	200 µL
Target:	Adenylate Kinase 2 (AK2)
Binding Specificity:	AA 147-183
Reactivity:	Human, Mouse, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Adenylate Kinase 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS)

Product Details

Immunogen:	This Zebrafish ak2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 147-183 amino acids from the Central region of Zebrafish ak2.
Clone:	RB53214
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	Adenylate Kinase 2 (AK2)
Alternative Name:	ak2 (AK2 Products)

Target Details


Background:	Catalyzes the reversible transfer of the terminal phosphate group between ATP and AMP. Plays an important role in cellular energy homeostasis and in adenine nucleotide metabolism. Adenylate kinase activity is critical for regulation of the phosphate utilization and the AMP de novo biosynthesis pathways. Plays a key role in hematopoiesis.
Molecular Weight:	26616
UniProt:	Q1L8L9
Pathways:	Nucleotide Phosphorylation , Ribonucleoside Biosynthetic Process

Application Details

Application Notes:	WB: 1:2000. IHC-P: 1:25. IHC-P: 1:25. FC: 1:25
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Expiry Date:	6 months




Flow Cytometry

Image 1. Overlay histogram showing ZF4 cells stained with (ABIN6242156 and ABIN6578282) (green line). The cells were fixed with 2 % paraformaldehyde (10 min) and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ((ABIN6242156 and ABIN6578282), 1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Goat Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/400 dilution for 40 min at 37 °C. Isotype control antibody (blue line) was rabbit IgG (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



Western Blotting

Image 2. All lanes : Anti-Zebrafish ak2 Antibody (Center) at 1:2000 dilution Lane 1: Zebrafish lysates Lane 2: ZF4 whole cell lysates Lane 3: human heart lysates Lane 4: mouse liver lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. (ABIN6242156 and ABIN6578282) staining Zebrafish ak2 in zebra fish body tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hours at 37 °C. A undiluted biotinylated goat polyvalent antibody was used as the

secondary antibody.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6242156.