antibodies -online.com







anti-INTS3 antibody (AA 1000-1034)



Image



| \sim | |
|---------|-------|
| ()\/\ | rview |
| \circ | |

| Quantity: | 200 μL |
|----------------------|--|
| Target: | INTS3 |
| Binding Specificity: | AA 1000-1034 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This INTS3 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | This INTS3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1000-1034 amino acids from human INTS3. |
| | |
| Clone: | RB56745 |
| Isotype: | Ig Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Target Details | |
| Target: | INTS3 |
| Alternative Name: | INTS3 (INTS3 Products) |
| Background: | Component of the Integrator complex. The Integrator complex is involved in the small nuclear |

Target Details


| | RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator |
|----------------------|---|
| | complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit |
| | (POLR2A) and is recruited to the U1 and U2 snRNAs genes. |
| Molecular Weight: | 118070 |
| UniProt [.] | O68F01 |

Application Details

| Application Notes: | WB: 1:2000 |
|--------------------|-----------------------|
| Restrictions: | For Research Use only |
| | |
| Handling | |

| Format: | Liquid |
|--------------------|--|
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Expiry Date: | 6 months |

Images



Western Blotting

Image 1. All lanes: Anti-INTS3 Antibody (C-Term) at 1:2000 dilution Lane 1: 293 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 118 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.