
antibodies -online.com







anti-HRH4 antibody (C-Term)



Image



Go to Product page

\sim			
	$ \backslash / \cap$	r\/I	$\triangle V$

Quantity:	400 μL
Target:	HRH4
Binding Specificity:	AA 359-390, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HRH4 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This HRH4 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 359-390 amino acids from the C-terminal region of human HRH4.
	populae between 605 656 animo acido nom the citeminarregion of hamaninin.
Clone:	RB1073
Clone: Isotype:	
	RB1073
Isotype:	RB1073 Ig Fraction
Isotype:	RB1073 Ig Fraction This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by
Isotype: Purification:	RB1073 Ig Fraction This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

Target Details

Background:	Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like	
	cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4.	
	Histamine receptor H4 belongs to the family 1 of G protein-coupled receptors and has the	
	highest homology to histamine receptor H3 among known G protein-coupled receptors.	
Molecular Weight:	44470	
UniProt:	Q96LD9	


Application Details

Application Notes:	WB: 1:2000. WB: 1:1000	
Restrictions:	For Research Use only	

Handling

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Expiry Date:	6 months	

Images



Western Blotting

Image 1. Anti-HRH4 Antibody (C-term) at 1:2000 dilution + human heart lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.