



[Go to Product page](#)

Datasheet for ABIN6242437
anti-MELK antibody (C-Term)

4 Images

Overview

Quantity:	400 µL
Target:	MELK
Binding Specificity:	AA 434-468, C-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MELK antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	This mouse Melk antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 434-468 amino acids from the C-terminal region of mouse Melk.
Clone:	RB51204
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	MELK
Alternative Name:	Melk (MELK Products)
Background:	Serine/threonine-protein kinase involved in various processes such as cell cycle regulation, self-

Target Details

renewal of stem cells, apoptosis and splicing regulation. Has a broad substrate specificity, phosphorylates BCL2L14, CDC25B, MAP3K5/ASK1 and ZNF622. Acts as an activator of apoptosis by phosphorylating and activating MAP3K5/ASK1. Acts as a regulator of cell cycle, notably by mediating phosphorylation of CDC25B, promoting localization of CDC25B to the centrosome and the spindle poles during mitosis. Plays a key role in cell proliferation. Required for proliferation of embryonic and postnatal multipotent neural progenitors. Phosphorylates and inhibits BCL2L14. Also involved in the inhibition of spliceosome assembly during mitosis by phosphorylating ZNF622, thereby contributing to its redirection to the nucleus. May also play a role in primitive hematopoiesis.

Molecular Weight: 72729

UniProt: [Q61846](#)

Application Details

Application Notes: WB: 1:1000. WB: 1:2000. WB: 1:1000. FC: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C, -20 °C

Expiry Date: 6 months




Flow Cytometry

Image 1. Overlay histogram showing MCF-7 cells stained with (ABIN6242437 and ABIN6577977) (red line). The cells were fixed with 2 % paraformaldehyde (10 min) and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ((ABIN6242437 and ABIN6577977), 1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (1583138) at 1/400 dilution for 40 min at 37 °C. Isotype control antibody (blue line) was rabbit IgG1 (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.



Western Blotting

Image 2. Anti-Melk Antibody (C-term) at 1:1000 dilution + mouse ovary lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 73 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.



Western Blotting

Image 3. Anti-Melk Antibody (C-term) at 1:2000 dilution + mouse spleen lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 73 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6242437.