

Datasheet for ABIN6242493
anti-TRKB antibody



[Go to Product page](#)

3 Images

Overview

Quantity:	400 µL
Target:	TRKB (NTRK2)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TRKB antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS)

Product Details

Immunogen:	This NTRK2 antibody is generated from a mouse immunized with a recombinant protein.
Clone:	1446CT494-85-83-49
Isotype:	IgG1 kappa
Purification:	This antibody is purified through a protein G column, followed by dialysis against PBS.

Target Details

Target:	TRKB (NTRK2)
Alternative Name:	NTRK2 (NTRK2 Products)
Background:	Receptor tyrosine kinase involved in the development and the maturation of the central and the peripheral nervous systems through regulation of neuron survival, proliferation, migration, differentiation, and synapse formation and plasticity. Receptor for BDNF/brain-derived

Target Details

neurotrophic factor and NTF4/neurotrophin- 4. Alternatively can also bind NTF3/neurotrophin-3 which is less efficient in activating the receptor but regulates neuron survival through NTRK2. Upon ligand-binding, undergoes homodimerization, autophosphorylation and activation. Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades. Through SHC1, FRS2, SH2B1, SH2B2 activates the GRB2-Ras-MAPK cascade that regulates for instance neuronal differentiation including neurite outgrowth. Through the same effectors controls the Ras-PI3 kinase-AKT1 signaling cascade that mainly regulates growth and survival. Through PLCG1 and the downstream protein kinase C-regulated pathways controls synaptic plasticity. Thereby, plays a role in learning and memory by regulating both short term synaptic function and long-term potentiation. PLCG1 also leads to NF-Kappa-B activation and the transcription of genes involved in cell survival. Hence, it is able to suppress anoikis, the apoptosis resulting from loss of cell-matrix interactions. May also play a role in neurotrophin- dependent calcium signaling in glial cells and mediate communication between neurons and glia.

Molecular Weight: 91999

UniProt: [Q16620](#)

Pathways: [RTK Signaling](#), [Neurotrophin Signaling Pathway](#), [cAMP Metabolic Process](#), [Skeletal Muscle Fiber Development](#), [Feeding Behaviour](#), [Dicarboxylic Acid Transport](#)

Application Details

Application Notes: WB: 1:1000. IHC-P: 1:25. FC: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Purified monoclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-20 °C

Expiry Date: 6 months



Flow Cytometry

Image 1. Overlay histogram showing SH-SY5Y cells stained with (ABIN6242493 and ABIN6577096) (green line). The cells were fixed with 2 % paraformaldehyde (10 min) and then permeabilized with 90 % methanol for 10 min. The cells were then incubated in 2 % bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ((ABIN6242493 and ABIN6577096), 1:25 dilution) for 60 min at 37 °C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(NA168821)) at 1/400 dilution for 40 min at 37 °C. Isotype control antibody (blue line) was mouse IgG1 (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



Western Blotting

Image 2. Western blot analysis of lysate from human brain tissue lysate, using NTRK2 Antibody (ABIN6242493 and ABIN6577096). (ABIN6242493 and ABIN6577096) was diluted at 1:1000. A goat anti-mouse IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20 µg.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. (ABIN6242493 and ABIN6577096) staining NTRK2 in human brain sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hour at 37 °C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.