.-online.com antibodies

Datasheet for ABIN6242500 anti-TR4 antibody (Ser439)

2 Images



Overview

Quantity:	200 μL
Target:	TR4 (NR2C2)
Binding Specificity:	AA 410-444, Ser439
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TR4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

Product Details

Immunogen:	This TAK1 (Ser439) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 410-444 amino acids from human TAK1.
Clone:	RB56649
Isotype:	Ig Fraction
Predicted Reactivity:	Rat
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	TR4 (NR2C2)
Alternative Name:	TAK1 (NR2C2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6242500 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Background:	Serine/threonine kinase which acts as an essential component of the MAP kinase signal
	transduction pathway. Plays an important role in the cascades of cellular responses evoked by
	changes in the environment. Mediates signal transduction of TRAF6, various cytokines
	including interleukin-1 (IL-1), transforming growth factor-beta (TGFB), TGFB-related factors like
	BMP2 and BMP4, toll-like receptors (TLR), tumor necrosis factor receptor CD40 and B-cell
	receptor (BCR). Ceramides are also able to activate MAP3K7/TAK1. Once activated, acts as an
	upstream activator of the MKK/JNK signal transduction cascade and the p38 MAPK signal
	transduction cascade through the phosphorylation and activation of several MAP kinase
	kinases like MAP2K1/MEK1, MAP2K3/MKK3, MAP2K6/MKK6 and MAP2K7/MKK7. These
	MAP2Ks in turn activate p38 MAPKs, c-jun N-terminal kinases (JNKs) and I-kappa-B kinase
	complex (IKK). Both p38 MAPK and JNK pathways control the transcription factors activator
	protein-1 (AP-1), while nuclear factor-kappa B is activated by IKK. MAP3K7 activates also IKBKB
	and MAPK8/JNK1 in response to TRAF6 signaling and mediates BMP2- induced apoptosis. In
	osmotic stress signaling, plays a major role in the activation of MAPK8/JNK1, but not that of
	NF-kappa-B. Promotes TRIM5 capsid-specific restriction activity.
Molecular Weight:	67196

UniProt:	O43318
Pathways:	TCR Signaling, Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling
	Pathway, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector
	Process, Production of Molecular Mediator of Immune Response, Tube Formation, Toll-Like
	Receptors Cascades

Application Details

Application Notes:	IF: 1:25. WB: 1:1000-1:2000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN6242500 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Handling

Storage:


Expiry Date:


4 °C,-20 °C

6 months

:

Images





Immunofluorescence

Image 1. Immunofluorescent analysis of 4 % paraforldehyde-fixed, 0.1 % Triton X-100 permeabilized HeLa (hun cervical epithelial adenocarcino cell line) cells labeling P3K7 with (ABIN6242500 and ABIN6578801) at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence ige showing cytoplasm and weak nucleus staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red). The nuclear counter stain is DI (blue).

Western Blotting

Image 2. All lanes : Anti-TAK1 (Ser439) Antibody at 1:1000-1:2000 dilution Lane 1: 293 whole cell lysate Lane 2: A431 whole cell lysate Lane 3: Raji whole cell lysate Lane 4: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 67 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6242500 | 09/10/2023 | Copyright antibodies-online. All rights reserved.