

Datasheet for ABIN6242506  
**anti-OR11H4 antibody (C-Term)**[Go to Product page](#)

## 1 Image

## Overview

|                      |                                       |
|----------------------|---------------------------------------|
| Quantity:            | 80 µL                                 |
| Target:              | OR11H4                                |
| Binding Specificity: | AA 239-271, C-Term                    |
| Reactivity:          | Human                                 |
| Host:                | Rabbit                                |
| Clonality:           | Polyclonal                            |
| Conjugate:           | This OR11H4 antibody is un-conjugated |
| Application:         | Western Blotting (WB)                 |

## Product Details

|               |   |
|---------------|---|
| Immunogen:    | This OR11H4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 239-271 amino acids from the C-terminal region of human OR11H4. |
| Clone:        | RB49843   |
| Isotype:      | Ig Fraction   |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification.  |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | OR11H4                                     |
| Alternative Name: | OR11H4 ( <a href="#">OR11H4 Products</a> ) |
| Background:       | Odorant receptor.                          |

## Target Details

Molecular Weight: 36886

UniProt: [Q8NGC9](#)

## Application Details

Application Notes: WB: 1:2000

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Expiry Date: 6 months

## Images



### Western Blotting

**Image 1.** Anti-OR11H4 Antibody (C-term) at 1:2000 dilution + MCF-7 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa. Blocking/Dilution buffer: 5 % NFDM/TBST.