

Datasheet for ABIN6242519
anti-PDXK antibody (AA 1-312)[Go to Product page](#)

2 Images

Overview

Quantity:	200 µL
Target:	PDXK
Binding Specificity:	AA 1-312
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PDXK antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This PDXK antibody is generated from a mouse immunized with a recombinant protein between 1-312 amino acids from human PDXK.
Clone:	1007CT2-1-1
Isotype:	IgG1 kappa
Purification:	This antibody is purified through a protein G column, followed by dialysis against PBS.

Target Details

Target:	PDXK
Alternative Name:	PDXK (PDXK Products)
Background:	Required for synthesis of pyridoxal-5-phosphate from vitamin B6.

Target Details

Molecular Weight: 35102

UniProt: [O00764](#)

Application Details

Application Notes: WB: 1:4000. IHC-P: 1:25

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified monoclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C


Expiry Date: 6 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. (ABIN6242519 and ABIN6577251) staining PDXK in human testis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hours at 37 °C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Western Blotting

Image 2. All lanes : Anti-PDXK Antibody (N-term) at 1:4000 dilution Lane 1: 293 whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: U-251 MG whole cell lysate Lane 4: Human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.